

Mini review: parasitoids of rice pests and prospects for enhancing biological control.

Red denotes added by jcatindig

Purple denotes added by jacheng – has also added names in Chinese of parasitoids and places but I could not reliably cut and paste them into this document

Summary of published information on insect parasitoids of rice pests.

TABLE 1. ORDER: HYMENOPTERA

Species name	Distribution Country/ region	Hosts	Notes (parasitism rates, alternative hosts, time of year, other information)	Reference (publication in English unless indicated otherwise)
<p>Family: Dryinidae Nymphal-adult parasites and predators, forelegs adapted for holding prey for feeding and ovipositing, on average feed on 3.2 nymphs and parasitize 4-9 nymphs per day, the parasite is brought into an early crop mainly via a parasitised host but also by migration of adults, parasitism increases with crop age but declines quickly when crop is mature. Highest parasitism rates 35-40%, September-October in dryland fields, affecting <i>Nephotettix virescens</i> and <i>S. furcifera</i> most (Chandra, 1980) This family parasitise nymphs or adults. They develop within a sac on the host's abdomen until ready to pupate. They emerge and pupate in a spun cocoon attached to the rice leaf or other surface (Chiu, 1979). Parasitism of <i>N. lugens</i> by dryinid wasps very low under 1-2%, (Kitamura, 1987) Relatively low total parasitism rates by dryinids ((<i>Echthrodolphax fairchildii</i>, <i>Haplogonatopus</i> spp. and <i>Pseudogonatopus</i> spp.) of <i>N. lugens</i> and <i>S. furcifera</i>: Wet season = 9.7%, dry season = 6.4% (Peña & Shepard, 1986) Parasitism of <i>L. striatellus</i> by dryinid wasps generally low under 10%, highest in August/September (Kitamura, 1987) Parasitism of <i>S. furcifera</i> by dryinid wasps was about 10%, most of time but up to 20% in June/July (Kitamura, 1987)</p>				
<p><i>Dicondylus indianus</i> Olmi</p> <p>(Dryinus)</p>	<p>Queensland, Australia Also India, Taiwan and the Philippines (citing Olmi, 1984)</p>	<p><i>Nilaparvata lugens</i> (Stål)</p>	<p>Needs to feed on host before ovipositing in them (obligate host-feeder/synovigenic). The adult parasitoids feed on first four instars and then lay eggs in all five nymph stages and adults – feeding and oviposition are non-concurrent as feeding usually causes the host to die. Females capture and restrain the host using their chelate foretarsi. Hosts are dropped off the plant if they have been fed upon or placed on their food plant if they have had eggs laid in them. Feeding on the hosts haemolymph is also important for survival. When host density high:</p>	<p>Sahragard et al, 1991</p> <p>n.b. For more on synovigeny and pro-ovigeny in relation to parasitoids - see Jervis & Kidd In: Theoretical Approaches to Biological Control, Bradford & Cornell (eds) (In CSU Orange library) and Jervis et al</p>
<p>Pseudonym of <i>Pseudogonatopus flavifemur</i></p>				

Comment [CSU1]: I don't understand this addition. I cannot find anything to suggest this is a pseudonym or anything under the name *Dryinus indianus* searching Web of Science, CAB abstracts and google

			<ul style="list-style-type: none"> the period of intense egg laying was short – peak egg laying reached sooner females lived longer average daily oviposition rate greater Regardless of host density an individual parasitoid did not lay more than 587 eggs	(2001) and Heimpel & Collier 1996 (in 'not used' collection of papers)
		<i>Nilaparvata lugens</i>	Nymph and adult parasite	CAB International, 2005
		<i>Nilaparvata lugens</i>	Nymph and adult parasitoid	Shepard et al, 2000
	Tropical Asia	<i>Laodelphax striatellus</i>	Nymph and adult	Reissig et al, 1986
<i>Digonatopus javanus</i> Keiffer		<i>Nilaparvata lugens</i>		CAB International, 2005
<i>Echthrodelpfax bicolor</i> Esaki & Hashimoto	Japan (citing Esaki & Hashimoto, 1936)	<i>Nilaparvata lugens</i>		Chiu, 1979
		<i>Nilaparvata lugens</i>	Nymph and adult parasite	CAB International, 2005
	Shimane, Japan	<i>Sogatella furcifera</i>		Kitamura, 1987 (Japanese – English abstract (have full text))
	Tropical Asia	<i>Nilaparvata lugens</i>	Nymph and adult parasitoid	Reissig et al, 1986
	Taiwan	<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Chu & Hirashima, 1981
<i>Echthrodelpfax fairchildii</i> Perkins*	India	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Nymph and adult parasite	Rhandhawa et al, 2006
		<i>Nilaparvata lugens</i>	Nymph and adult parasite	CAB International, 2005
	Madhya Pradesh, India	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> (Horvath)		Yadav & Pawar, 1989
	Japan	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>Laodelphax striatellus</i> Fallen	Daily fecundity about 15-25 eggs	Ito & Yamada, 2007
		<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>		Barrion & Litsinger, 1994 p.184

		<i>Laodelphax striatellus</i> , <i>Perkinsiella saccharicida</i> , <i>Nephotettix spp.</i>		
	Japan	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Semi-solitary – daily fecundity approx 20 To oviposit the female holds the host with its mandibles and chelae and paralyzes it by stinging. The egg is oviposited under the wing bud on either side Mating required to produce females	Yamada & Ikawa, 2003
	Philippines	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Relatively low total parasitism rates by dryinids (<i>Echthrodelphax fairchildii</i> , <i>Haplogonatopus spp.</i> and <i>Pseudogonatopus spp.</i>) Wet season = 7.6%, dry season = 11.2%	Peña & Shepard, 1986
	Mandya (Karnataka), India	<i>Nilaparvata lugens</i>	Nymph and adult parasite	Manjunath et al, 1978
	Philippines		Form larval sac on the dorsal side of the host's thorax	Chandra, 1980
	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Less common	van Vreden & Ahmadzabidi, 1986
	Philippines	<i>Nilaparvata lugens</i>	Reared on BPH	Barrion et al, 1981
	India (citing Rai, pers. comm,)	<i>Nilaparvata lugens</i>		Chiu, 1979
			Hyperparasitised by <i>Cheiloneurus hawaiiensis</i> Perkins – Hawaii	Guerrieri & Viggiani, 2005
<i>Echthrodelphax spp.</i>	India, Japan, Korea, Philippines, Taiwan	<i>Sogatella furcifera</i> <i>Nilaparvata lugens</i> <i>Nephotettix cincticeps</i> <i>Nephotettix virescens</i>		Greathead, 1982
<i>Gonatopus yasumatsui</i> Olmi		<i>Nilaparvata lugens</i>		Barrion & Litsinger, 1994 p.186
	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Less common	van Vreden & Ahmadzabidi, 1986
<i>Gontpopus sp.</i>			Hyperparasitised by <i>Cheiloneurus exitiosus</i> Perkins – Australia, Samoa, Fiji, Guam, India, Malaysia, Philippines And <i>Cheiloneurus pachycephalus</i> Perkins – Ohio, USA	Guerrieri & Viggiani, 2005
<i>Haplogonatopus sp. nr. americanus</i>	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Less common	van Vreden & Ahmadzabidi, 1986

<i>Haplogonatopus apicalis</i> Perkins*	India	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Nymph and adult parasite	Rhandhawa et al, 2006
	Madhya Pradesh, India	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>		Yadav & Pawar, 1989
		<i>Nilaparvata lugens</i>	“During larval development, it feeds on the host’s body fluid. It protrudes from the abdomen of its host as a black to grayish sac. After 7 to 10 days, the larval sac splits to expose a whitish larva. Pupation occurs outside the host’s body.”	http://zj.shuidao.cn/IRRI/beneficials/Scientific_name_Haplogonatopus_apicalis_Perkins_78.htm
		<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>Laodelphax striatellus</i>		Barrion & Litsinger, 1994 p.186
		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
	Japan	<i>Sogatella furcifera</i>	The sex ratio of progeny was associated with the age of adult parasitoid, the development of the host (male ratio lowest in third instars) and the sex of the host but not the host density. Oviposition frequency was associated with the development of the host with highest occurring in 3 rd -instar nymphs Those ovipositing in females were larger than those ovipositing in males	Kitamura & Iwami, 1998 (Japanese – abstract in English)
	Tropical Asia	<i>Sogatella furcifera</i>	Nymph and adult parasitoid	Reissig et al, 1986
<i>Haplogonatopus atratus</i> Esaki & Hashimoto		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
		<i>Laodelphax striatellus</i>		Barrion & Litsinger, 1994 p.188
	Japan	<i>Laodelphax striatellus</i>	Host specific Mating required to produce females	Yamada & Kawamura, 1999
		<i>Laodelphax striatellus</i>	Appears to maximise its reproductive capacity by being fecund and laying as many eggs in as many hosts as possible rather than guarding the eggs it has already laid	Yamada & Kitashiro, 2002
	Sinan County, Guizhou, China	<i>Laodelphax striatellus</i>	48.2% of total parasitism of delphacids (abstract not explicit about species) by 6 parasitoid species	Chen, 1989 (in Chinese – English abstract)

	Japan	<i>Laodelphax striatellus</i> <i>Sogatella furcifera</i>	The most dominant parasitoid species Predator as well as a parasitoid Observed in the field that the parasite mainly affects <i>L. Striatellus</i> In the field although no difference was observed in the lab, 54.9% mortality caused by combined parasitic and predatory action of the parasitoid (mostly due to parasitisation) in the lab – thought to be higher than in the field	Kitamura, 1982
	Shimane, Japan	<i>Laodelphax striatellus</i>		Kitamura, 1987 (Japanese – English abstract (have full text))
	Japan	<i>Laodelphax striatellus</i>	Solitary parasitoid	Yamada & Miyamoto, 1998
	Tropical Asia	<i>Laodelphax striatellus</i>	Nymph and adult parasitoid	Reissig et al, 1986
<i>Haplogonatopus japonicus</i> Esaki & Hashimoto	Japan (citing Esaki & Hashimoto, 1931; Sakai, 1932; Esaki & Mochizucki, 1941)	<i>Nilaparvata lugens</i>		Chiu, 1979
Alternative spellings: <i>H. japonica</i> , <i>H. japonicas</i> Chen, 1989 and Zhang & Jin, 1992 appear to indicate <i>H. japonicas</i> is synonymous with <i>H. apicalis</i>	Sinan County, Guizhou, China	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	36.5% of total parasitism of delphacids (abstract not explicit about species) by 6 parasitoid species	Chen, 1989 (in Chinese – English abstract)
	Guangdong Province, China	<i>Nilaparvata lugens</i>	A common parasitoid species found in that area	Ying, 1982
	Zhejiang, China	<i>Sogatella furcifera</i>	Appears to indicate this is a synonym of <i>H. apicalis</i> Malathion, dipterex (trichlorfon) and MTMC (melolcarb) – highly toxic to parasitoid Methamidophos, chlordimeform hydrochloride and dimethoate – mildly toxic to parasitoid	Zhang & Jin, 1992 (Chinese – English abstract)
	China	<i>Sogatella furcifera</i>		Huang, 1994; Chen 1990 (in Chinese – English abstracts)
	Shimane, Japan	<i>Sogatella furcifera</i>		Kitamura, 1987 (Japanese – English abstract (have full text))

<i>Haplogonatopus oratorius</i> Westwood		<i>Laodelphax striatellus</i> ,		Barrion & Litsinger, 1994 p.188
<i>Haplogonatopus orientalis</i> Rohwer	India	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Nymph and adult parasite	Rhandhawa et al, 2006
	India	<i>Nilaparvata lugens</i>	<i>N. lugens</i> has a symbiotic relationship with <i>Candida</i> sp. The yeast load declined considerably when parasitised by this parasitoid	Shankar & Baskaran, 1992 (abstract only)
		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	CAB International, 2005
	Sri Lanka	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	The age-advanced larva forms an outstanding sac on the dorsal part of the abdomen of the host. 40% parasitism rates reached – but was not persistent and failed to control the population of planthoppers	Ôtake et al, 1976
	India	<i>Nilaparvata lugens</i>	Identified from parasitized nymphs and adults collected from the insectary and rice fields. Egg parasitoid – symbiotic yeast (important for the planthoppers nutrition) decrease as parasites grow	Shankar & Baskaran, 1988
<i>Haplogonatopus</i> sp./spp.	Philippines	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>		Peña & Shepard, 1986
	Mandya (Karnataka), India	<i>Nilaparvata lugens</i>	Nymph and adult parasite	Manjunath et al, 1978
	Philippines	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> Nymphs of <i>Nephotettix virescens</i>	Form a larval sac on the dorsal side of the host's abdomen of <i>N. lugens</i> and <i>S.furcifera</i>	Chandra, 1980
	Philippines	<i>Nilaparvata lugens</i>	Reared on BPH	Barrion et al, 1981
	India (citing Rai, pers. comm.), Sri Lanka (citing Santa et al, unpubl.)	<i>Nilaparvata lugens</i>		Chiu, 1979
	Tropical Asia	<i>Nilaparvata lugens</i>	Nymph and adult parasitoid	Reissig et al, 1986
	Fiji, India, Japan, Korea, Philippines	<i>Sogatella furcifera</i> <i>Nilaparvata lugens</i> <i>Nephotettix virescens</i>		Greathead, 1982

			Hyperparasitised by <i>Cheiloneurus exitiosis</i> Perkins – Australia, Samoa, Fiji, Guam, India, Malaysia, Philippines	Guerrieri & Viggiani, 2005
	Malaysia	<i>Sogatella furcifera</i> <i>Nilaparvata lugens</i>	Claims <i>S. furcifera</i> and <i>N. lugens</i> were the two main planthoppers causing much damage in Malaysia. Suggests little was known about parasitoids of planthoppers at that time	Ooi, 1982
<i>Monogonatopus orientalis</i> Rohwer	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Abundant	van Vreden & Ahmadzabidi, 1986
<i>Monogonatopus</i> sp.	Taiwan	<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Chu & Hirashima, 1981
<i>Pseudogonatopus otakei</i> Olmi	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Less common	van Vreden & Ahmadzabidi, 1986
<i>Pseudogonatopus atratus</i> Esaki & Hashimoto		<i>Nilaparvata lugens</i>		
<i>Pseudogonatopus flavifemur</i> Esaki & Hashimoto * Alternative spelling: <i>P. flavifemur</i> should be <i>Dicondylus indianus</i> Olmi)		<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>Nephotettix</i> spp.		Barrion & Litsinger, 1994 p.188
	Shimane, Japan	<i>Nilaparvata lugens</i>		Kitamura, 1987 (Japanese – English abstract (have full text))
	Japan (citing Esaki & Hashimoto, 1933; 1936; Esaki, 1932; Sakai, 1932)	<i>Nilaparvata lugens</i>		Chiu, 1979
	Sinan County, Guizhou, China	<i>Nilaparvata lugens</i>	10.8% of total parasitism of delphacids (abstract not explicit about species) by 6 parasitoid species	Chen, 1989 (in Chinese – English abstract)
	Philippines	<i>Nilaparvata lugens</i>	4.7% parasitism rate	Dayanan & Esteban, 1996 (abstract only)
	Philippines	<i>Nilaparvata lugens</i>	Dryinids feed on hosts killing them and so fewer healthy hosts are available for parasitisation. In their experiments a female parasitoid fed on up to ten third-instar hosts. They suggest superparasitism is mainly an artefact of laboratory experiments because of the small experimental area and the long time span. Observed	Chua et al, 1984

Comment [CSU2]: I cannot find anything for this species searching Web of Science, Google and CAB abstracts

			<p>superparasitism only once in the field. Females laid fewer eggs when another female was present. Increasing parasitoid density increases 'mutual interference' between them – the area searched was reduced and hosts were handled for longer. In biological control programs host and parasitoid populations stabilise as high searching efficiency reduces the average host density and parasitoids are attracted to the most dense populations where 'mutual interference' reduces oviposition. In addition, mutual interference encourages dispersal of the parasitoid after introduction. Normally in the field parasitism rates are low - <20%</p>	
	Philippines	<i>Nilaparvata lugens</i>	Nymphs and adults parasite	CAB International, 2005
	Philippines			Chandra, 1980
	Philippines	<i>Nilaparvata lugens</i>	Reared on BPH	Barrion et al, 1981
	Taiwan	<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Chu & Hirashima, 1981
	Philippines (Los Baños, Cabanatuan, Bayombing, Kiangan and Banaue)		Not specific about which parasites were affecting which planthoppers or leafhoppers (N. lugens and S.furcifera found in all locations along with Nephrotettix virescens, Nephrotettix nigropictus and Recilia dorsalis)	Heong et al, 1992
			Hyperparasitised by <i>Cheiloneurus exitiosus</i> Perkins – Australia, Samoa, Fiji, Guam, India, Malaysia, Philippines	Guerrieri & Viggiani, 2005
<i>Pseudogonatopus flavifemur</i> Esaki & Hashimoto	Philippines			Chandra, 1980
<i>Pseudogonatopus fulgori</i> Nakagawa		<i>Laodelphax striatellus</i> <i>Sogatella furcifera</i>		Barrion & Litsinger, 1994 p.188
	Japan	<i>Laodelphax striatellus</i> <i>Sogatella furcifera</i>	<p>In <i>L. striatellus</i> it took on average 26.1 days for eggs to develop to pupae at 24°C In <i>S. furcifera</i> it took on average 25.9 days for eggs to develop to pupae at 24°C Eggs failed to develop in <i>N. lugens</i> Development temperature threshold: 13.8°C Development from egg to pupa took from 16 days at</p>	Kitamura, 1989 (Japanese – English abstract)

Comment [CSU3]:

flavifemur and flavifemur both listed by Chandra – doesn't look like a typo - searching Web of Science, Google and CA abstracts

			32°C to 42.5 days at 20°C	
	Shimane, Japan	<i>Laodelphax striatellus</i> <i>Sogatella furcifera</i>		Kitamura, 1987 (Japanese – English abstract (have full text))
<i>Pseudogonatopus hospes</i> Perkins*	Madhya Pradesh, India Also found in Japan and Malaysia (citing Anonymous, 1978 and Chiu, 1979)	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>		Yadav & Pawar, 1989
	Hawaii - Kauai; Oahu; Molokai; Maui; Hawaii	<i>Nilaparvata lugens</i>	Purposely Introduced	http://www2.bishopmuseum.org/HBS/checklist/species.asp?grp=A&rthropod&taxID=-1208574862
		<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>		Barrion & Litsinger, 1994 p.191
	Thailand (citing Napompeth, unpublished)	<i>Nilaparvata lugens</i>		Chiu, 1979
		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Less common	van Vreden & Ahmadzabidi, 1986
	Hawaii	Sugar cane leafhopper	Introduced from China	Swezey, 1925
			Hyperparasitised by <i>Cheiloneurus exitiosus</i> Perkins – Australia, Samoa, Fiji, Guam, India, Malaysia, Philippines	Guerrieri & Viggiani, 2005
<i>Pseudogonatopus nudus</i> Perkins*		<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>		Barrion & Litsinger, 1994 p.188
	Philippines, India, Sri Lanka, Thailand, China, Taiwan, Indonesia and Malaysia	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Unable to distinguish between <i>P. sarawaki</i> and <i>P. nudus</i>	Olmi 1991-2
	Philippines	<i>Nilaparvata lugens</i>	3.3% parasitism rate	Dayanan & Esteban, 1996 (abstract only)
	Philippines	<i>Nilaparvata lugens</i>	Dominant species of dryinid parasitoid of <i>N. lugens</i> in Philippines (citing Chandra, 1980)	Chua et al, 1984

		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
	Philippines	<i>Nilaparvata lugens</i>	Reared on BPH	Barrion et al, 1981
	Tropical Asia	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Nymph and adult parasitoid	Reissig et al, 1986
		<i>Nilaparvata lugens</i>	Nymph and adult parasitoid	Shepard et al, 2000
		<i>Nilaparvata lugens</i>	Dominant species on this host	Greathead, 1982
<i>Pseudogonatopus ponomarenkoi</i> Moczar	Tropical Asia	<i>Sogatella furcifera</i>	Nymph and adult parasitoid	Reissig et al, 1986
<i>Pseudogonatopus</i> nr. <i>pusanus</i> Olmi	Madhya Pradesh, India Also found in Japan and Malaysia (citing Anonymous, 1978 and Chiu, 1979)	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>		Yadav & Pawar, 1989
<i>Pseudogonatopus</i> <i>pusanus</i> Olmi		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
<i>Pseudogonatopus</i> <i>sarawaki</i> Moczar*		<i>Nilaparvata lugens</i> <i>Sogatella</i> sp.		Barrion & Litsinger, 1994 p.188
		<i>Nilaparvata lugens</i>	Unable to distinguish between <i>P. sarawaki</i> and <i>P. nudus</i>	Olmi 1992
		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
	Tropical Asia	<i>Nilaparvata lugens</i>	Nymph and adult parasitoid	Reissig et al, 1986
<i>Pseudogonatopus</i> sp./ spp.	Philippines	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>		Peña & Shepard, 1986
	Philippines	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	From larval sac on dorsolateral side of the host's abdomen	Chandra, 1980
		<i>Nilaparvata lugens</i>	Common parasitoid	Ooi & Shepard, 1994
	Philippines (Los Baños, Cabanatuan, Bayombing, Kiangan and Banaue)		Not specific about which parasites were affecting which planthoppers or leafhoppers (<i>N. lugens</i> and <i>S. furcifera</i> found in all locations along with <i>Nephotettix virescens</i> , <i>Nephotettix nigropictus</i> and <i>Recilia dorsalis</i>)	Heong et al, 1992
	Japan, Philippines, Taiwan	<i>Sogatella furcifera</i> <i>Nilaparvata lugens</i> <i>Nephotettix cincticeps</i>		Greathead, 1982

		<i>Nephotettix virescens</i>		
			Hyperparasitised by <i>Cheiloneurus gonatopdis</i> Perkins – Australia, Madagascar and Mauritius	Guerrieri & Viggiani, 2005
	Taiwan	<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Chu & Hirashima, 1981
<i>Pseudozonatus hospes</i> Perkins	Malaysia	<i>Sogatella furcifera</i> <i>Nilaparvata lugens</i>	Claims <i>S. furcifera</i> and <i>N. lugens</i> were the two main planthoppers causing much damage in Malaysia. Suggests little was known about parasitoids of planthoppers at that time	Ooi, 1982
Family: Encyrtidae hyperparasites of Dryinidae (Chiu 1979)				
<i>Cheilonerurus exitiosus</i> Perkins		<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Also a hyperparasite of other parasitoids (<i>Gonatopus</i> sp., <i>Haplogpnatopus</i> sp., <i>Pseudogonatopus hospes</i> , <i>P. flavifemur</i>)	Guerrieri & Viggiani, 2005
<i>Chrysopophagus australiae</i> Perkins	Solomon Islands (citing MacQuillan, 1974)	<i>Nilaparvata lugens</i>		Chiu, 1979
<i>Echthrogonatopus exitiosus</i> Perkins	Solomon Islands (citing MacQuillan, 1974)	<i>Nilaparvata lugens</i>		Chiu, 1979
		<i>Nilaparvata lugens</i>	Nymph and adult parasite	CAB International, 2005
Family: Eulophidae				
<i>Ootetrastichus nr beatus</i>	Fiji (citing Hinckley, 1963)	<i>Nilaparvata lugens</i>	Egg parasite Parasitism appeared to be rare.	Chiu, 1979
		<i>Nilaparvata lugens</i>	Egg parasite	CAB International, 2005
	Hawaii	Sugar cane leafhopper	Introduced from Fiji	Sweezey, 1925
	South Africa	<i>Numicia viridis</i>	More important in control of this host than <i>Oligosita</i> sp. because it is more “robust”	Dick & Thompson, 1969
	South Africa and Swaziland	<i>Numicia viridis</i>	Associated with sugar cane and wild grasses – along with <i>Oligosita</i> sp. effectively controls the pest	Charleston et al, 2003
<i>Ootetrastichus nr. formosanus</i> <i>Timberlake</i>	Philippines	<i>Nilaparvata lugens</i>	Reared on BPH	Barrion et al, 1981
	Tropical Asia	<i>Nilaparvata lugens</i>	Egg parasitoid	Reissig et al, 1986
	Hawaii	Sugar cane leafhopper	Introduced from Formosa	Sweezey, 1925
<i>Ootetrastichus</i> spp.	Philippines (Los Baños, Cabanatuan, Bayombing, Kiangnan		Not specific about which parasites were affecting which planthoppers or leafhoppers (<i>N. lugens</i> and <i>S.furcifera</i> found in all locations along with <i>Nephotettix virescens</i> ,	Heong et al, 1992

	and Banaue)		Nephotettix nigropictus and Recilia dorsalis)	
<i>Tetrastichus formosanus</i> Timberlake		<i>Nilaparvata lugens</i>	Egg parasite	CAB International, 2005
	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Common parasitoid	van Vreden & Ahmadzabidi, 1986
	Tropical Asia	<i>Nilaparvata lugens</i>	Egg parasitoid	Reissig et al, 1986
	Philippines (Los Baños, Cabanatuan, Bayombing, Kiangan and Banaue)		Not specific about which parasites were affecting which planthoppers or leafhoppers (N. lugens and S. furcifera found in all locations along with Nephotettix virescens, Nephotettix nigropictus and Recilia dorsalis)	Heong et al, 1992
	Thailand	<i>Nilaparvata lugens</i>		Wongsiri et al, 1980
Family: Mymaridae				
Egg parasites – quick to migrate into crops from alternative hosts in other habitats, quickly establish cause rapid mortality – therefore important as biocontrol agents (Chandra, 1980)				
Most myramid parasites around in May and June (parasitism rates 11.3-29.6%) and again in September to November (3.3 – 38.1%) (Chiu, 1979)				
<i>Anagrus sp. nr flaveolus</i> Waterhouse	Sri Lanka	<i>Nilaparvata lugens</i>		Fowler et al, 1991
	Japan (citing Otake 1970a,b); 1976a,b)	<i>Nilaparvata lugens</i>	Egg parasite Parasitism rates 44.5% (in Zentuz) and 66.9% (in Kagawa) and influences pest populations in early growth stages (citing Otake, 1976b) Parasitism could be easily detected through the transparent chorion of the host egg when the parasite larva was at least half grown.	Chiu, 1979
	Japan	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>Laodelphax striatellus</i>	No preference shown between <i>N. lugens</i> , <i>S. furcifera</i> and <i>L. striatellus</i> Have a strong tendency to disperse – important for overwintering away from paddy fields in non-delphacid species Outlines methods for determining parasitism rates	Ôtake, 1977
			Unfertilised eggs all males (citing Otake, 1969) – see also <i>Anagrus sp.</i> entry outlines difficulties associated with and methods for sampling egg parasitoid populations in the field and estimating parasitism rates	Chandra, 1980
	Philippines	<i>Nilaparvata lugens</i>	Reared on BPH	Barrion et al, 1981

		<i>Nilaparvata lugens</i>	Reared the parasitoid in the laboratory by designing a new rearing technique	Chandra & Dyck, 1987
	Taiwan	<i>Nilaparvata lugens</i>	Egg parasitoid	Ya u-i & Hirashima, 1981
		<i>Nilaparvata lugens</i>	Wasp parasitizes 15-30 eggs a day	Shepard et al, 2000
	Japan	<i>Laodelphax striatellus</i>	An overwhelmingly dominant species attacking eggs of <i>Laodelphax striatellus</i> on cultivated plots.	Otake, 1970
	India	<i>Sogatella furcifera</i>	Parasitization was more on resistant genotype with five times more than with susceptible variety.	Nalini, 2005.
	Tropical Asia	<i>Sogatella furcifera</i>	Egg parasitoid	Reissig et al, 1986
<i>Anagrus flaveolus</i> Waterhouse			Unfertilised eggs all females – see also <i>Anagrus</i> sp. entry	Chandra, 1980
	China	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>Tagosodes pusanus</i>	Dominant species Had significant preference of <i>N. lugens</i> and <i>S. furcifera</i> when on rice – but on grass preferred <i>T. pusanus</i> Non-rice habitats are important for the conservation of the rice planthopper parasitoids	Yu et al, 1998 (Chinese – English abstract)
	China	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	In rice fields and adjacent habitats Population dynamics (sex ratio, body size and parasitoid growth rate) were influenced by: host eggs, host plants of parasitoids and the surrounding habitat. The parasitoids obtained nutrients from nectar and pollen of the non-rice flowers.	Yu et al, 1996 (Chinese – English Abstract)
	China	<i>Toya</i> spp. and <i>Tagosodes pusanus</i>	Important egg parasitoid of delphacids in rice and non-rice habitats Habitat dominated by grasses close to paddy fields may act as a reservoir of parasitoids of rice planthoppers	Yu 1996 (Chinese – English Abstract)
	Japan (citing Yasumatsu & Watanabe, 1965)	<i>Nilaparvata lugens</i>		Chiu, 1979
	San Miguel de Tucumán, Argentina	<i>Delphacodes kuscheli</i>		Liljeström & Virla, 2004
	India	<i>Nilaparvata lugens</i>	Effective parasitoid of <i>Empoasca</i> sp. and also parasitised <i>Amrasca b. biguttula</i> Activity peaked in July and October-November but active throughout year – paper does not mention <i>N.</i>	Singh et al, 1993 (abstract only)

			<i>lugens</i>	
	Saxian county, Fujian province	Planthoppers	Abstract does not identify the host/s involved <i>Anagrus flaveolus</i> and two closely related species are main natural control agents of rice planthoppers Graminaceous weeds important for overwintering natural enemies	Lo & Zhou, 1980 (Chinese – English Abstract)
	Japan	<i>Laodelphax striatellus</i>	66%-96% parasitism measured by 'trap method' - late June to early August	Hachiya, 1995 (Japanese – English abstract)
	Malaysia	<i>Sogatella furcifera</i>	<i>Anagrus</i> spp.: • turn host eggs orange or yellow orange and parasitoids can be seen through the transparent chorion 47% maximum parasitism	Watanabe et al, 1992
	India	<i>Sogatella furcifera</i>	Egg parasite	Rhandhawa et al, 2006
	Taiwan	<i>Nephotettix cincticeps</i> <i>Nilaparvata lugens</i>	Egg parasite	Chu & Hirashima, 1981
<i>Anagrus frequens</i>		<i>Nilaparvata lugens</i>		CAB International, 2005
Synonyms: <i>Anagrus armatus</i> , <i>A. cicadulinae</i>	Malaysia	<i>Sogatella furcifera</i>	<i>Anagrus</i> spp.: • turn host eggs orange or yellow orange and parasitoids can be seen through the transparent chorion 47% maximum parasitism	Watanabe et al, 1992
	India	<i>Sogatella furcifera</i>	Egg parasite	Rhandhawa et al, 2006
	Hawaii	Sugar cane leafhopper	Introduced from Australia	Swezey, 1925
	Queensland, Australia Hawaii, Fiji, Philippines, Taiwan	<i>Perkinsella saccharicida</i> (Fiji and Hawaii) <i>Kelisia sporobolicola</i> (Hawaii), <i>Perkinsella maidis</i> (Hawaii), corn leafhopper (Philippines), <i>Perkinsella</i> sp. (Taiwan)		Triapitsyn & Beardsley, 2000
<i>Anagrus incarnatus</i> Haliday	Piacenza province, Italy	<i>Nilaparvata lugens</i>		Chiappini et al, 1999
	Taiwan, Britain,	<i>Nilaparvata lugens</i>		Chen & Yu, 1989

	Denmark, Belgium, Japan, Korea, Bangladesh			
	China	<i>Nilaparvata lugens</i>		Chiappini et al, 1998
	Japan	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>N. muiri</i> and <i>N. bakeri</i> (citing Chantarasa-ard, 1984).	<i>N. lugens</i> and <i>S. furcifera</i> do not overwinter in Japan but spread from overseas each year. <i>A. incarnatus</i> capable of overwintering in eggs of <i>N. muiri</i>	Chantarasa-ard, 1984
	Japan	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>Laodelphax striatellus</i> <i>N. bakeri</i> Muir <i>N. muiri</i> <i>Harmalia albicolli</i> Motchulsky, <i>Sogatella longifurcifera</i> Esaki et Ishihara, <i>S. panicola</i> Ishihara, <i>Terthron albovittatum</i> Matsumura, <i>Zuleica nipponica</i> Matsumura et Ishihara, <i>Nephotettix cincticeps</i> Uhler <i>Macrosteles orientalis</i> Vilbaste	Does not have a preference between <i>S. furcifera</i> , <i>N. lugens</i> and <i>L. Striatellus</i> (70%+ parasitism rates) Can attack eggs at any stage of development. The parasitism was higher and the parasite more abundant in the field in September compared to August. Parasitism rates for alternative hosts: <i>N. bakeri</i> Muir (71.4% in lab), <i>N. muiri</i> (39.9% in lab; 32.4% in field), <i>Harmalia albicolli</i> Motchulsky (93.8% in lab; 0% in field), <i>Sogatella longifurcifera</i> Esaki et Ishihara (65.8% in lab; 26% in field), <i>S. panicola</i> Ishihara (17.8% in lab; 18.1% in field), <i>Terthron albovittatum</i> Matsumura (40% in lab; 64.7% in field), <i>Zuleica nipponica</i> Matsumura et Ishihara (8.8% in field) <i>Nephotettix cincticeps</i> Uhler (5.3% in lab) and <i>Macrosteles orientalis</i> Vilbaste (63% in lab).	Chantarasa-ard & Hirashima, 1984
	Japan	<i>Sogatella furcifera</i>	Most dominant egg parasitoid	Chantarasa-ard et al, 1984
	Navarra, Spain England (citing Enoch, 1914), Belgium (citing Debauche, 1948; citing Mathot, 1969), Holland and Austria (citing Soyka, 1946; 1955), Bulgaria, Yugoslavia and Greece (citing	<i>Cicadella viridis</i> <i>Delphacodes fairmairei</i> and <i>Juncus effuses</i> (citing Trjapitsyn, 1997) other Jassidae, Delphacidae and Odonata (citing Bakkendorf, 1926; Whalley, 1956)		Baquero & Jordano, 1999

	Donev, 1978; 1985a; 1985b), Japan, Korea and Bangladesh (citing Shad & Hiroshima, 1984), Poland and Turkmenistan (citing Trjapitsyn, 1997)			
<i>Anagrus longithbulosus</i>			Proposes the synonymy of <i>A. longitubulosus</i> under <i>A. (paranagrus) perforator</i>	Triapitsyn, 2001(abstract only)
Possibly should be <i>Anagrus longitubulosus</i>	China	<i>Nilaparvata lugens</i>	Found to be one of the dominant species in parasitoid communities in terms of numbers and importance as a control agent. All species in the parasitoid community, including the dominant species, fluctuated. Numbers were related to the numbers of the host and stage of rice growth. Parasitism rates in relation to rice growing period: early 76%, middle 70% and late 50%.	Mao et al, 2002b (Chinese – English Abstract)
<i>Anagrus nilaparvatae</i> Pang et Wang		<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>Laodelphax striatellus</i>	Prefers <i>N. lugens</i> over <i>S. furcifera</i> - better survival rates and fecundity when emerging from <i>N. lugens</i>	Lou & Cheng, 2001
	China	<i>Nilaparvata lugens</i>	The parasitoid's host preference is influenced by HIPVs	Xiang et al, 2008
	China	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>Laodelphax striatellus</i> <i>Sogatella vibix</i> (Haupt), <i>Soatella longifurcifera</i> (Esaki and Ishihara), <i>Nilaparvata bakeri</i> (Muir), <i>Nilaparvata muiri</i> China, <i>Toya propinqua</i> (Feiber) and <i>Toya tuberculosa</i> (Distant)	Alternatives hosts are used by the parasite in winter when <i>N.lugens</i> is scarce (Citing Wand and Pang, 1986) Most common before early August mainly attacking <i>L. striatella</i> and <i>S. furcifera</i> (citing Xu and Cheng, 1988) Ideal temp: 27.41°C – fecundity and survival of immatures greatly reduced at high temperatures (citing Cheng & Xu, 1991)	Chiappini et al, 1998
	China	<i>Nilaparvata lugens</i>	Both <i>N. lugens</i> and <i>A. nilaparvata</i> more attracted to plants treated with jasmonic acid (JA) than to control plants. Treatment with JA also enhanced parasitism	Lou et al, 2005a
		<i>Nilaparvata lugens</i>	Plants had to be damaged by <i>N. lugens</i> feeding to be	Lou et al, 2005b

Comment [CSU4]: I cannot find anything searching Web of Science, CAB abstracts and google – assumed it should be *Anagrus longitubulosus*

			<p>more attractive to <i>A. nilaparvata</i> – females were not attracted to volatiles from undamaged hosts plants or those just mechanically damaged.</p> <p>Parasitoids were attracted most to intermediate host densities. Very high or low densities of host densities did not attract the parasitoids.</p>	
		<i>Nilaparvata lugens</i>	<p>Demonstrates ethylene signalling is involved in induction of plant volatiles by herbivory by <i>N. lugens</i>. Ethylene production by the plant is involved in regulating the amount and mix of induced volatiles (citing Huang et al, 2005).</p>	Lu et al, 2006
South China		<i>Nilaparvata lugens</i>	<p>Parasitoid communities were steadier in IPM areas compared to non-IPM areas (presumably using conventional pesticide regimes).</p> <p>Temperature, species, number of species, distance from species pool, the host species present and the control regime employed all influenced the reestablishment and maintenance of the parasitoid communities.</p>	Mao et al, 2002a
China		<i>Nilaparvata lugens</i>	<p>Found to be one of the dominant species in parasitoid communities in terms of numbers and importance as a control agent.</p> <p>All species in the parasitoid community, including the dominant species, fluctuated. Numbers were related to the numbers of the host and stage of rice growth.</p> <p>Parasitism rates in relation to rice growing period: early 76%, middle 70% and late 50%.</p>	Mao et al, 2002b
Guangdong, China		<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>Toya tuberculosa,</i> <i>Nilaparvata bakeri,</i> <i>Sogatella panicola</i> [<i>S. vibix</i>], <i>T. propinqua</i>	<p>A dominant parasitoid species – peaking in October and June</p> <p>Parasitism rate of <i>N. lugens</i>: 20-60%</p> <p>Parasitism rate of <i>S. furcifera</i>: 40.3-92.6%</p>	Li & He, 1991 (in Chinese – English abstract)
		<i>Sogatella furcifera</i>	<p>Displayed no preference for eggs of differing age but took longer to develop and body size and fecundity of the adult wasp decreased in older eggs. No effect was observed on sex ratio and emergence rate. Fecundity was associated with body size.</p>	Zhu et al, 1993 (Chinese – English Abstract)

			Preferred <i>N. lugens</i> eggs over <i>S. furcifera</i> – in which bigger, more fecund wasps were produced – but development was slower.	
<p><i>Anagrus optabilis</i> Perkins</p> <p>Mao et al, 2002a seems to say that <i>A. paranilaparvatae</i> is a pseudonym of <i>A. optabilis</i></p> <p>Triapitsyn, 2001 (abstract only), proposes the synonymy of <i>A. paranilaparvatae</i> under <i>A. optabilis</i></p> <p>Synonyms: <i>Paranagrus optabilis</i> Perkins, <i>Paranagrus osborni</i> Fullway, <i>Anagrus panicolae</i> Sahad (Triapitsyn & Beardsley, 2000)</p>	India	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Egg parasite	Rhandhawa et al, 2006
	Andhra Pradesh and Tamil Nadu India Taiwan	<i>Nilaparvata lugens</i>		CAB International, 2005
	India	<i>Nilaparvata lugens</i>	<i>N. lugens</i> has a symbiotic relationship with <i>Candida</i> sp. The yeast disappears completely when parasitised by this particular parasitoid	Shankar & Baskaran, 1992 (abstract only)
	China	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	In rice fields and adjacent habitats Population dynamics (sex ratio, body size and parasitoid growth rate) were influenced by: host eggs, host plants of parasitoids and the surrounding habitat. The parasitoids obtained nutrients from nectar and pollen of the non-rice flowers.	Yu et al, 1996 (Chinese – English Abstract)
	China	<i>Toya</i> spp. and <i>Tagosodes pusanus</i>	Important egg parasitoid of delphacids in rice and non-rice habitats Habitat dominated by grasses close to paddy fields may act as a reservoir of parasitoids of rice planthoppers	Yu 1996 (Chinese – English Abstract)
	Thailand (citing Yasumatsu et al, 1975; Nishida et al, 1976)	<i>Nilaparvata lugens</i>	Egg parasite Important species - More abundant than <i>Paracentrobia yasumatsui</i> or a few other unidentified parasites. Parasitism rates <10% to 100% (citing Nishida et al, 1976)	Chiu, 1979
	Doi Saket, Lamphun, San Pa Tong, Chom Thong, Hot and Chai Nat, Thailand	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Parasitism rates varied from 100% in Doi Saket to 14.2% in Chai Nat	Hiroshima, 1979
	Malaysia	<i>Nilaparvata lugens</i>	<i>Anagrus</i> spp.: • turn host eggs orange or yellow orange and parasitoids can be seen through the transparent chorion • 20-60% parasitism rate	Watanabe et al, 1992
	Sri Lanka	<i>Nilaparvata lugens</i>	Parasitism rates of <i>Anagrus</i> spp. (<i>optabilis</i> and <i>flaveolus</i>): site B = 10%	Fowler et al, 1991

			Parasitism rates did not appear to be dependent on batch size and was also unrelated to host egg density at the tiller level but significantly positively related to the number of eggs per plant.	
Tropical Asia	<i>Nilaparvata lugens</i>	Egg parasitoid		Reissig et al, 1986
	<i>Nilaparvata lugens</i>	Wasp parasitizes 15-30 eggs a day		Shepard et al, 2000
Peninsular Malaysia	<i>Nilaparvata lugens</i>	Abundant		van Vreden & Ahmadzabidi, 1986
Taiwan	<i>Nilaparvata lugens</i> <i>Laodelphax striatellus</i> <i>Sogatella furcifera</i>	<i>N. lugens</i> : 33.88% parasitism rate for egg masses and 19.56% for individual eggs from 9 localities <i>L. striatellus</i> : 45.90% parasitism rate for egg masses and 30.54% for individual eggs from 14 localities <i>S. furcifera</i> : 20-40% parasitism rate from 7 localities		Miura et al, 1981
India	<i>Nilaparvata lugens</i>	Parasitized eggs collected from the insectary and rice fields. Egg parasitoid – symbiotic yeast (important for the planthoppers nutrition) decrease as parasites grow		Shankar & Baskaran, 1988
Japan	<i>Nilaparvata lugens</i> <i>Laodelphax striatellus</i> <i>Sogatella furcifera</i>	<i>N. lugens</i> : 86.7% parasitism rate <i>L. striatellus</i> : 70.5% parasitism <i>S. furcifera</i> : 79.6% parasitism rate		Sahad , 1984
Thailand	<i>Sogatella furcifera</i>	Important egg parasitoid		Miura et al, 1979
Navarra, Spain Queensland, Australia (citing Perkins, 1905), Hawaii, Japan, Taiwan, Thailand and Bangladesh (citing Sahad & Hirashima, 1984), Malaysia, India, Sri lank, New Guinea, Indonesia, Mauritius, Philippines, Fiji, Samoa and Guam (citing Trjapitsyn, 1997), Ecuador (citing De Santis & Fidalgo, 1994),	<i>Nilaparvata lugens</i> <i>Nilaparvata muii</i> <i>Laodelphax striatellus</i> <i>Sogatella furcifera</i> <i>Saccharosdyne procerus</i> <i>Zulieca nipponica</i> and <i>Leersia japonica</i> (citing Sahda & Hirashima) <i>Perkinsella saccharicida</i> <i>Perkinsella thompsoni</i> and <i>Pundaloya simplicia</i> (citing Chiappini, et al 1996)			Baquero & Jordana, 1999

	South Africa (citing Triapitsyn, 1997)			
	Thailand	<i>Nilaparvata lugens</i> <i>Nephotettix</i> spp.		Wongsiri et al, 1980
	Queensland, Australia Hawaii, Philippines, Taiwan, Indonesia	<i>Perkinsella saccharicida</i> (Australia and Hawaii), <i>Kelisia emoloa</i> (Hawaii), leafhoppers (Indonesia), <i>Perkinsella vastatrix</i> , corn leafhopper (Philippines), <i>Perkinsella</i> sp. (Taiwan)		Triapitsyn & Beardsley, 2000
	Malaysia	<i>Sogatella furcifera</i> <i>Nilaparvata lugens</i>	Claims <i>S. furcifera</i> and <i>N. lugens</i> were the two main planthoppers causing much damage in Malaysia. Suggests little was known about parasitoids of planthoppers at that time	Ooi, 1982
<i>Anagrus paranilaparvatae</i> Pang & Wang	China	<i>Nilaparvata lugens</i>	Dominant species after mid-August mainly parasitising <i>N. Lugens</i> (citing Xu and Cheng, 1988) Ideal temp: 31.87°C - highest mortality at 18°C (citing Cheng & Xu, 1991)	Chiappini, 1998
Triapitsyn, 2001 (abstract only), proposes the synonymy of <i>A. paranilaparvatae</i> under <i>A. optabilis</i>	South China	<i>Nilaparvata lugens</i>	Parasitoid communities were steadier in IPM areas compared to non-IPM areas (presumably using conventional pesticide regimes). Temperature, species, number of species, distance from species pool, the host species present and the control regime employed all influenced the reestablishment and maintenance of the parasitoid communities. Seems to say that <i>A. paranilaparvatae</i> is a pseudonym of <i>A. optabilis</i>	Mao et al, 2002a
	China	<i>Nilaparvata lugens</i>	Found to be one of the dominant species in parasitoid communities in terms of numbers and importance as a control agent. All species in the parasitoid community, including the dominant species, fluctuated. Numbers were related to the numbers of the host and stage of rice growth. Parasitism rates in relation to rice growing period: early 76%, middle 70% and late 50%.	Mao et al, 2002b

	Jiangsu, China	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>Laodelphax striatellus</i>	Showed a preference for eggs of <i>N. lugens</i> (I think this is in relation to <i>S. furcifera</i> and <i>L. striatella</i> but maybe also to <i>N. lugens</i> in other parts of the plant – not clear from abstract) located in the upper and middle parts of the plant and in tiller tissue.	Hu, 1992 (Chinese – English Abstract)
	Guangdong, China	<i>Nilaparvata lugens</i>		Li & He, 1991(Chinese – English Abstract)
<i>Anagrus perforator</i> Perkins	India	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Egg parasite	Rhandhawa et al, 2006
			Proposes the synonymy of <i>A. longitubulous</i> under <i>A. (paranagrus) perforator</i>	Triapitsyn, 2001(abstract only)
Synonym: <i>Paranagrus perforator</i>	Andhra Pradesh, India	<i>Nilaparvata lugens</i>	Egg parasite	CAB International, 2005
	Malaysia	<i>Sogatella furcifera</i>	<i>Anagrus</i> spp.: • turn host eggs orange or yellow orange and parasitoids can be seen through the transparent chorion • 47% maximum parasitism	Watanabe et al, 1992
	China	<i>Sogatella furcifera</i>		Chiappini, 1998
	China	<i>Nilaparvata lugens</i>	Found to be one of the dominant species in parasitoid communities in terms of numbers and importance as a control agent. All species in the parasitoid community, including the dominant species, fluctuated. Numbers were related to the numbers of the host and stage of rice growth. Parasitism rates in relation to rice growing period: early 76%, middle 70% and late 50%.	Mao et al, 2002b (Chinese – English Abstract)
	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Less common	van Vreden & Ahmadzabidi, 1986
	Fiji, Hawaii, Japan, Philippines	<i>Hirozuunka japonica</i> , (Japan), <i>Megamelus Proserpina</i> , <i>Nephotettix virescens</i> , and <i>Sogatella furcifera</i> (Philippines)		Triapitsyn & Beardsley, 2000
<i>Anagrus</i> sp./spp.	Guangdong Province, China	<i>Nilaparvata lugens</i>	Common parasitoids found in that area	Ying, 1982

	China	<i>Sogatella furcifera</i>	Main natural enemy in early rice fields – May-July	Luo & Zhou, 1986 (Chinese – English Abstract)
	Taiwan, Japan (citing Lin, 1974; Fukuda, 1934; Kuno, 1973) Fukuoka, Japan (citing Kuno & Hokyo, 1970) and Malaysia (citing Heong (pers comm..))	<i>Nilaparvata lugens</i>	Egg parasite – when the parasite larvae was half grown parasitism can easily seen through the host egg's transparent chorion (citing Otake 1970) Parasitism rates 10%-15% (citing Kuno & Hokyo, 1970)	Chiu, 1979
	Taipei and Pingtung, Taiwan (citing Lin, 1974)	<i>Nilaparvata lugens</i>	Most prevalent (93% in Taipei) of egg parasitoids among myramids and trichogrammatids (citing Lin, 1974)	Chui, 1979
	Philippines	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>Nephotettix virescens</i> (Distant)	Most common genera of mymarid parasitoid When adult lands on rice plant it walks quickly all over it, drumming on the surface with their antennae. When they find a host egg mass they drum more energetically close to the eggs. Oviposition occurs by the wasp first drilling through the leaf epidermis. The drumming appears to be involved in locating the eggs and finding a suitable place to drill. Failure rate is high: 95% attempts fail to penetrate and of those that do 89% do not successfully oviposit in an egg. When parasite density is high 1-3 eggs laid but only one will develop. The parasitised egg expands rapidly to several times the original size – hatch after 2 days, transparent, instar - moults at one day – turns yellowish which can then be seen within the host egg – pupates within 24hrs – pupa: bright orange-red turning brown, larvae wiggle lots within the egg to break up the host tissue for ingestion - pupation takes 6-7 days – adults emerge from host eggs 11-13 days after oviposition – males emerge first Parthenogenic (whether males or females emerge depends on the species – see also <i>A. flaveolus</i> and <i>A. nr. flaveolus</i> enteries) and gametogenetic reproduction Females lived 2-6 days in lab conditions – but did not oviposit after 3 days	Chandra, 1980

		Longevity did not affect fecundity because most of the eggs laid within 24hrs.	
Philippines	<i>Nilaparvata lugens</i>	Reared on BPH	Barrion et al, 1981
Indonesia	<i>Nilaparvata lugens</i>	Low to higher rates of parasitism in wet season and more uniform higher levels in dry seasons. Several species	Claridge et al, 1999
	<i>Nilaparvata lugens</i>	Common egg parasitoid	Ooi & Shepard, 1994
Vietnam	<i>Sogatella furcifera</i>	Egg parasitoid with 72.5% parasitism	Tao & Ngoan, 1970
Thailand	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Dominant egg parasite of <i>S.furcifera</i> in north and Central Plain and of <i>N. lugens</i> in the Central Plain (found not to attack leafhoppers)	Vungsilabutr, 1981
India		listed as an important natural enemy of planthoppers	Pasalu et al, 2004
		Studied the effect of Bt transgenic rice on the dispersal of planthoppers, leafhoppers and their egg parasitoid wasps. <i>Anagrus</i> spp. tended to disperse towards block of non- transgenic rice	Chen et al, 2003 (Chinese – English abstract)
India	<i>Nilaparvata lugens</i>	Along with <i>Oligosita</i> sp. – the most common parasitoids of <i>N. lugens</i>	Gupta & Pawar, 1989 (abstract only)
Philippines (Los Baños, Cabanatuan, Bayombing, Kiangan and Banaue)		Not specific about which parasites were affecting which planthoppers or leafhoppers (<i>N. lugens</i> and <i>S.furcifera</i> found in all locations along with <i>Nephotettix virescens</i> , <i>Nephotettix nigropictus</i> and <i>Recilia dorsalis</i>)	Heong et al, 1992
Japan, Fiji, Korea, Malaysia, Philippines, Sri Lanka, Thailand, Taiwan, Vietnam	<i>Sogatella furcifera</i> <i>Nilaparvata lugens</i> <i>Nephotettix cincticeps</i> <i>Nephotettix nigropicta</i> <i>Nephotettix virescens</i>	Dominant genus of Mymaridae in rice fields Which hosts are affected varies in different countries (see p.375)	Greathead, 1982
Malaysia	<i>Sogatella furcifera</i> <i>Nilaparvata lugens</i>	Claims <i>S. furcifera</i> and <i>N. lugens</i> were the two main planthoppers causing much damage in Malaysia. Suggests little was known about parasitoids of planthoppers at that time	Ooi, 1982
Taiwan	<i>Nephotettix cincticeps</i> <i>Nilaparvata lugens</i>	Egg parasite	Chu & Hirashima, 1981

	Zhejiang, China	<i>Sogatella furcifera</i>	The draining of rice fields at the 8-9 leaf stage of the first crop and 12-13 leaf stage of the 2 nd crop reduced number of eggs laid and hatching and therefore resulted in lower populations of the host and encouraged natural enemies - parasitism rates of <i>S. furcifera</i> by <i>Anaphes</i> sp. was 9.5% lower	Zhang, 1991 (Chinese, English abstract)
<i>Anaphes taprobanicum</i>		<i>Nilaparvata lugens</i>	Egg parasite	CAB International, 2005
<i>Anaphes</i> spp/ sp.	Solomon Islands (citing MacQuillan, 1974)	<i>Nilaparvata lugens</i>	Egg parasite	Chiu, 1979
	Taiwan	<i>Nephotettix cincticeps</i> <i>Nilaparvata lugens</i>	Egg parasite	Chu & Hirashima, 1981
<i>Gonatocerus</i> sp.	Thailand (citing Yasumatsu et al, 1975)	<i>Nilaparvata lugens</i>	Egg parasite	Chiu, 1979
	Korea (citing Yasumatsu, personal communication)	<i>Nilaparvata lugens</i>		Ôtake, 1977
	Philippines	Specific to <i>Nephotettix virescens</i>		Chandra, 1980
	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Less common	van Vreden & Ahmadzabidi, 1986
	Thailand	<i>Nephotettix virescens</i> <i>Nephotettix nigropictus</i>	Was found not to attack <i>N. lugens</i> or <i>S. furcifera</i>	Vungsilabutr, 1981
	India		listed as an important natural enemy of planthoppers	Pasalu et al, 2004
	Philippines (Los Baños, Cabanatuan, Bayombing, Kiangan and Banaue)		Not specific about which parasites were affecting which planthoppers or leafhoppers (<i>N. lugens</i> and <i>S. furcifera</i> found in all locations along with <i>Nephotettix virescens</i> , <i>Nephotettix nigropictus</i> and <i>Recilia dorsalis</i>)	Heong et al, 1992
	Thailand	<i>Nilaparvata lugens</i>		Wongsiri et al, 1980
	Korea, Philippines, Taiwan, Thailand	<i>Nephotettix cincticeps</i> <i>Nephotettix virescens</i>	Usually the most abundant mymarid egg parasitoid	Greathead, 1982
	Taiwan	<i>Nephotettix cincticeps</i> <i>Nilaparvata lugens</i>		Chu & Hirashima, 1981
<i>Lymaenon</i> sp.	Taiwan (citing Lin, 1974)	<i>Nilaparvata lugens</i>	Egg parasite	Chiu, 1979

<i>Lymaenon longicrus</i>	Saxian county, Fujian province	Planthoppers	Abstract does not identify the host/s involved Graminaceous weeds important for overwintering natural enemies	Lo & Zhou, 1980 (Chinese – English Abstract)
<i>Mymar? Indica Mani</i>	Taiwan (citing Loin, 1974 and Chiu et al, unpubl.)	<i>Nilaparvata lugens</i>	Egg parasite	Chiu, 1979
	India			Gupta & Poorani, 2008
	Taiwan	<i>Nephotettix cincticeps</i> <i>Nilaparvata lugens</i>		Chu & Hirashima, 1981
<i>Mymar taprobanicum Ward</i>	Thailand (citing , Yasumatsu et al 1975)	<i>Nilaparvata lugens</i>	Egg parasite Important species (citing Yasumatsu et al 1975)	Chiu, 1979
	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Less common	van Vreden & Ahmadzabidi, 1986
	India, East and Far East, Korea and probably many other countries		Suggests <i>M. indica</i> is synonymous with <i>M. taprobanicum</i>	Subba Rao, 1976
	Sri Lanka			Gupta & Poorani, 2008
	Thailand	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>		Wongsiri et al, 1980
<i>Mymar</i> sp.	Philippines	<i>Nilaparvata lugens</i>	Rare	Chandra, 1980
	Philippines	<i>Nilaparvata lugens</i>	Reared on BPH	Barrion et al, 1981
<i>Polynema</i> sp.	Thailand (citing , Yasumatsu et al 1975)	<i>Nilaparvata lugens</i>	Egg parasite	Chiu, 1979
Family: Pteromalidae				
<i>Panstenon</i> sp	Saxian county, Fujian province	Planthoppers	Abstract does not identify the host/s involved Graminaceous weeds important for overwintering natural enemies	Lo & Zhou, 1980 (Chinese – English Abstract)
Family: Scelionidae				
<i>Baeus</i> sp.	Mandya (Karnataka), India	<i>Nilaparvata lugens</i>	Egg parasite	Manjunath et al, 1978

Comment [CSU5]: This genus appears mainly to parasitise spiders

<i>Gryon</i> sp.	Mandya (Karnataka), India	<i>Nilaparvata lugens</i>	Egg parasite	Manjunath et al, 1978
<i>Oxyscella</i> sp.	Mandya (Karnataka), India	<i>Nilaparvata lugens</i>	Egg parasite	Manjunath et al, 1978
Family: Trichogrammatidae Egg parasites (Chandra, 1980)				
<i>Alphelinoidae</i> sp.	Taiwan (citing Fukada, 1934)	<i>Nilaparvata lugens</i>	Egg parasite	Chiu, 1979
	Taiwan		Egg parasitoid	Ya u-i & Hirashima, 1981
<i>Oligosita aesopi</i> Girault	Malaysia	<i>Sogatella furcifera</i>	<i>Oligosita</i> spp.: • turn host eggs dark yellow and chorion is dark grey and so parasitoids cannot be seen >5% parasitism rate	Watanabe et al, 1992
	China	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	In rice fields and adjacent habitats Population dynamics (sex ratio, body size and parasitoid growth rate) were influenced by: host eggs, host plants of parasitoids and the surrounding habitat. The parasitoids obtained nutrients from nectar and pollen of the non-rice flowers.	Yu et al, 1996 (Chinese – English Abstract)
	China	<i>Toya</i> spp. and <i>Tagosodes pusanus</i>	Important egg parasitoid of delphacids in rice and non- rice habitats Habitat dominated by grasses close to paddy fields may act as a reservoir of parasitoids of rice planthoppers	Yu 1996 (Chinese – English Abstract)
		<i>Nilaparvata lugens</i>	Parasitizes 2-8 eggs a day	Shepard et al, 2000
	Tropical Asia	<i>Sogatella furcifera</i>	Egg parasitoid	Reissig et al, 1986
	<i>Oligosita naias</i> Girault	Tamil Nadu, India	<i>Nilaparvata lugens</i>	Egg parasite
China		<i>Toya</i> spp. and <i>Tagosodes pusanus</i>	Important egg parasitoid of delphacids in rice and non- rice habitats Habitat dominated by grasses close to paddy fields may act as a reservoir of parasitoids of rice planthoppers	Yu 1996 (Chinese – English Abstract)
China		<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	In rice fields and adjacent habitats Population dynamics (sex ratio, body size and parasitoid	Yu et al, 1996 (Chinese – English Abstract)

			growth rate) were influenced by: host eggs, host plants of parasitoids and the surrounding habitat. The parasitoids obtained nutrients from nectar and pollen of the non-rice flowers.	
	Philippines; Australia-Queensland; India-Andhra Pradesh; India-Karnataka; India-Orissa; India-Tamil Nadu; Malaysia; Peoples' Republic of China-Zhejiang	<i>Nilaparvata lugens</i>		http://www.catalogueoflife.org/show_species_details.php?record_id=4533017
	Muda area, Malaysia	<i>Nilaparvata lugens</i>	<i>Oligosita</i> spp.: • turn host eggs dark yellow and chorion is dark grey and so parasitoids cannot be seen 34-68% parasitism rate	Watanabe et al, 1992
		<i>Nilaparvata lugens</i>	Wasp parasitizes 2-8 eggs a day.	Shepard et al, 2000
	India	<i>Sogatella furcifera</i>	Egg parasite	Rhandhawa et al, 2006
<i>Oligosita nepholettica</i> Mani	Saxian county, Fujian province	Planthoppers	Abstract does not identify the host/s involved Graminaceous weeds important for overwintering natural enemies	Lo & Zhou, 1980 (Chinese – English Abstract)
	Indonesia	<i>Nilaparvata lugens</i>		Claridge et al, 1999
	Taiwan	<i>Nephotettix cincticeps</i> <i>Nilaparvata lugens</i>	Listed in this paper as <i>Oilogsita nephotetticum</i> but almost certainly a ‘typo’ Egg parasite	Chu & Hirashima, 1981
<i>Oligosita Shibuyae</i>	Taiwan	<i>Nephotettix cincticeps</i> <i>Nilaparvata lugens</i>	Egg parasite	Chu & Hirashima, 1981
<i>Oligosita tachikawai</i>	Andhra Pradesh, India	<i>Nilaparvata lugens</i>	Egg parasite	CAB International, 2005
<i>Oligosita yasumatsuii</i> Viggiani et Subba Rao	Andhra Pradesh, India	<i>Nilaparvata lugens</i>	Egg parasite	CAB International, 2005
	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Abundant	van Vreden & Ahmadzabidi, 1986
		<i>Nilaparvata lugens</i>	Common egg parasitoid	Ooi & Shepard, 1994

	Tropical Asia	<i>Laodelphax striatellus</i>		Reissig et al, 1986
	Indonesia	<i>Nilaparvata lugens</i>		Claridge et al, 1999
	Thailand	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>		Wongsiri et al, 1980
<i>Oligosita</i> sp./spp.			Drumming the surface of the rice leaves and oviposition occurs in a similar manner to <i>Anagrus</i> spp. (see that section). Dissecting the host eggs is not a good method of determining parasitisation because larvae and pupae of the wasp is very delicate and easily destroyed. Larvae within the eggs are difficult to observe as they do not wiggle much. Hatched larvae are light yellow, becoming brighter when they pupate – 5 days after oviposition the pupae, with pink eyes and body segments, can be observed easily through the chorion. Parasitised eggs are discernable because they retain a lemon yellow colour with a black band at the base. Adults emerge 11-12 days after oviposition – males emerge first. Most of the females eggs laid on first day after emergence Mating results in all females Less active as less fecund than mymarids	Chandra, 1980
	Sri Lanka	<i>Nilaparvata lugens</i>	More abundant than the <i>Anagrus</i> spp. parasitising the host. Parasitism rates: site A = 18% and site B = 32.7% Parasitism rates did not appear to be dependent on batch size and was also unrelated to host egg density at the tiller level but significantly positively related to the number of host eggs per plant.	Fowler et al, 1991
	Mandya (Karnataka), India	<i>Nilaparvata lugens</i>	Egg parasite	Manjunath et al, 1978
	Thailand (citing Yasumatsu, 1975), Taiwan (citing Lin, 1974)	<i>Nilaparvata lugens</i>	Egg parasite Effective at suppressing pest populations (citing Yasumatsu, 1975)	Chiu, 1979
	Chiand Dao and Mae Theng, Thailand			Hiroshima, 1979
	Guangdong	<i>Nilaparvata lugens</i>		(no citation for this)

				addition)
	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Less abundant	van Vreden & Ahmadzabidi, 1986
	Philippines	<i>Nilaparvata lugens</i>	Reared on BPH	Barrion et al, 1981
	Indonesia	<i>Nilaparvata lugens</i>	50% reduction on egg hatching of BPH when plants were exposed to the activity of the parasitoids.	Claridge, 1996
	Indonesia	<i>Nilaparvata lugens</i>	Low to higher rates of parasitism in wet seasons and more uniform higher levels through dry seasons.	Claridge et al, 1999
	Thailand	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Two species of Oligosita were found, one only affected the leafhoppers (<i>N. virescens</i> and <i>N. nigropictus</i>) and the other only affected the planthoppers	Vungsilabutr, 1981
	India		listed as an important natural enemy of planthoppers	Pasalu et al, 2004
	India	<i>Nilaparvata lugens</i>	Along with <i>Anagrus</i> sp. – the most common parasitoids of <i>N. lugens</i>	Gupta & Pawar, 1989 (abstract only)
	Philippines (Los Baños, Cabanatuan, Bayombing, Kiangan and Banaue)		Not specific about which parasites were affecting which planthoppers or leafhoppers (<i>N. lugens</i> and <i>S. furcifera</i> found in all locations along with <i>Nephotettix virescens</i> , <i>Nephotettix nigropictus</i> and <i>Recilia dorsalis</i>)	Heong et al, 1992
	In all SE Asia areas Fiji, India, Korea, Malaysia, Philippines, Solomon Is., Sri Lanka, Thailand, Taiwan	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i> <i>Nephotettix cinciteps</i> <i>Nephotettix nigropicta</i> <i>Nephotettix virescens</i>	Of equal importance to <i>Anagrus</i> spp. and dominates where there is multiparasitism and can be more abundant (citing IRRI, 2978)	Greathead, 1982
	South Africa	<i>Numicia viridis</i>		Dick & Thompson, 1969
	Malaysia	<i>Sogatella furcifera</i> <i>Nilaparvata lugens</i>	Claims <i>S. furcifera</i> and <i>N. lugens</i> were the two main planthoppers causing much damage in Malaysia. Suggests little was known about parasitoids of planthoppers at that time	Ooi, 1982
	Taiwan	<i>Nephotettix cinciteps</i> <i>Nilaparvata lugens</i>	Egg parasite	Chu & Hirashima, 1981
	South Africa and Swaziland	<i>Numicia viridis</i>	Associated with sugar cane and wild grasses	Charleston et al, 2003
<i>Paracentrobia andoi</i> Ishii	Taiwan and Japan (citing Suenaga; Lin, 1974)	<i>Nilaparvata lugens</i>	Egg parasite Parasitism rate extremely low	Chiu, 1979
Synonym: <i>P.</i>		<i>Nilaparvata lugens</i>	Egg parasite	CAB International,

<i>japania</i> (Chu & Hirashima, 1981)				2005
	Saxian county, Fujian province	Planthoppers	Abstract does not identify the host/s involved Gramineaceous weeds important for overwintering natural enemies	Lo & Zhou, 1980 (Chinese – English Abstract)
	Taiwan	<i>Nilaparvata lugens</i>	The female of the parasitoid attacked almost all the eggs when she came in contact with an egg mass of the hopper.	Miura et al, 1979
	Tropical Asia	<i>Nilaparvata lugens</i>	Egg parasitoid	Reissig et al, 1986
		<i>Nephotettix cincticeps</i>	Under lab conditions found that the eggs of <i>N. lugens</i> , <i>S. furcifera</i> and <i>L. striatellus</i> were not parasitised	Vungsilabutr et al, 1996 (abstract only)
	Japan and Taiwan		Replaces <i>Oligosita</i> spp.	
	Taiwan	<i>Nephotettix cincticeps</i> <i>Nephotettix nigropictus</i> <i>Nephotettix virescens</i> <i>Nilaparvata lugens</i>	Egg parasite	Chu & Hirashima, 1981
<i>Paracentrobia garuda</i> Subba Rao	Thailand (citing Yasumatsu et al, 1975)	<i>Nilaparvata lugens</i>	Egg parasite Effective at suppressing pest populations (citing Yasumatsu, 1975)	Chiu, 1979
		<i>Nilaparvata lugens</i>	Egg parasite	CAB International, 2005
	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Less common	Van Vreden & Ahmadzabidi, 1986
	Thailand	<i>Nilaparvata lugens</i>		Wongsiri et al, 1980
	Thailand	<i>Nephotettix virescens</i> <i>Nephotettix nigropictus</i>	Was found not to attack <i>N. lugens</i> or <i>S. furcifera</i>	Vungsilabutr, 1981
<i>Paracentrobia yasumatsui</i> Subba Rao	Thailand (citing Yasumatsu et al, 1975)	<i>Nilaparvata lugens</i>	Egg parasite Effective at suppressing pest populations (citing Yasumatsu, 1975)	Chiu, 1979
		<i>Nilaparvata lugens</i>	Egg parasite	CAB International, 2005
	Peninsular Malaysia	<i>Nilaparvata lugens</i>	Less common	Van Vreden & Ahmadzabidi, 1986
	Thailand	<i>Nilaparvata lugens</i>		Wongsiri et al, 1980
<i>Stephanodes</i> sp.	Philippines	<i>Nilaparvata lugens</i>	Reared on BPH	Barrion et al, 1981
<i>Trichogramma</i> sp.	Taiwan (citing Fukuda, 1934)	<i>Nilaparvata lugens</i>	Egg parasite	Chiu, 1979

*Major parasites of *N. Lugens* according to Ooi & Shepard (1994)

TABLE 2. ORDER: STREPSIPTERA

Nymphal-adult parasites (Chandra, 1980)

Hosts are not killed quickly but the parasitoid develops within the host allowing it to continue to damage the crop (Chandra, 1980)

Favoured wetland environment – parasitism higher in rainy seasons – but mostly below 10% - higher rates rare and only occurred in *N. lugens* (Chandra, 1980)

Species name	Distribution Country/ region	Hosts	Notes (parasitism rates, alternative hosts, time of year, other information)	Reference (publication in English unless indicated otherwise)
Family: Elenchidae				
<i>Elenchus japonicus</i> Esaki & Hashimoto Alternative spelling: <i>E. japonicus</i>	Japan (citing Esaki & Hashimoto, 1932; Esaki, 1932; Sakai, 1932; Okada, 1973; Kuno 1973)	<i>Nilaparvata lugens</i>		Chiu, 1979
	Guangdong Province, China	<i>Nilaparvata lugens</i>	A common parasitoid species found in that area	Ying, 1982
	Japan	<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
	Shimane, Japan	<i>Nilaparvata lugens</i> <i>Laodelphax striatellus</i> <i>Sogatella furcifera</i>	Parasitism rate of delphacids (predominantly <i>S. furcifera</i>): 0.1-26.7%, starting in July and highest in August	Kitamura, 1987 (Japanese – English abstract (have full text))
	India	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Nymph and adult parasite	Rhandhawa et al, 2006
<i>Elenchus koebelei</i> Pierce	Fiji (citing Hinckley, 1963)	<i>Nilaparvata lugens</i>	Occasionally parasitised <i>N. lugens</i>	Chiu, 1979
		<i>Nilaparvata lugens</i>	Nymph and adult parasite	CAB International, 2005
<i>Elenchus yasumatsui</i> Kefune & Hirashima*	Philippines	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	In <i>N. lugens</i> : parasitism rates of <i>N. lugens</i> relatively low in wet (1.4%) and dry season (5.5%) In <i>S. furcifera</i> : highest parasitism rate of 26% in March Overall parasitism rates relatively low in wet (7.4%) and dry seasons (9.8%)	Peña & Shepard, 1986
	Philippines	<i>Nilaparvata lugens</i>	10% parasitisation rate	Dayanan & Esteban, 1996 (abstract only)
		<i>Nilaparvata lugens</i>		Ooi & Shepard, 1994
	Thailand (citing Kifune	<i>Nilaparvata lugens</i>		

Comment [CSU6]: This genus appears to be very important as control agents

	& Hirashima, 1975; Otake, 1976)			
		<i>Nilaparvata lugens</i>	Females wingless, life-span 1 to 2 months, do not emerge fully from host (just the head outside) Males winged, lifespan 1-2 days, emerge and seek out females Somehow they mate and produce 500 to 2000 triungulins or larvae They do not oviposit in the host but the tiny C-shape triungulins out the host and bore inside	http://www.ctpm.uq.edu.au/software/riceipm/keys/Html/Elenchus.htm
	Sarawak	<i>Nilaparvata lugens</i>		Hirashima, 1978
	Thailand and Sarawak	<i>Sogatella furcifera</i>		Hiroshima, 1979
	northern Thailand (citing Yasumatsu et al, 1975)	<i>Nilaparvata lugens</i>	Plays a significant in control of <i>N. lugens</i> (citing Yasumatsu et al, 1975) Parasitism rates 30% - 90% (citing FAO, 1975)	Chiu, 1979
	Philippines	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>	Viviparous – females produce 1000-2000 triungulins – 0.15mm long, light yellow, slightly curved, well developed eyes, legs and caudal setae – crawl and jump – in lab most die within an hour – enter host by piercing intersegmental membranes - shrink and transform into cylindrical legless larvae – 7 instars – males pupate with anterior end poking out of the hosts abdomen – female pupates within the host – adult males emerge out of the host and mate with adult females via the cephalothorax which is exposed. Parasitised host's have: smaller genitalia, an extended abdomen and discoloured bodies as well as having the male parasitoids extruding from their abdomens and the females cephalothorax visible. Host and females adults die soon after triungulins have emerged. Hosts vacated by males are vulnerable to disease via the hole left. Showed preference for <i>N. lugens</i> over <i>S. furcifera</i> in the field	Chandra, 1980
	Tropical Asia	<i>Laodelphax striatellus</i>	Nymph and adult parasitoid	Reissig et al, 1986
	Thailand	<i>Nilaparvata lugens</i> <i>Sogatella furcifera</i>		Wongsiri et al, 1980

<i>Elenchus sp/spp.</i>	Sri Lanka (citing Santa et al., unpublished)	<i>Nilaparvata lugens</i>	Hosts are rendered sterile (citing Kuno, 1973)	Chiu, 1979
	India	<i>Nilaparvata lugens</i>	<i>N. lugens</i> has a symbiotic relationship with <i>Candida</i> sp. The yeast load declined considerably when parasitised by this parasitoid	Shankar & Baskaran, 1992 (abstract only)
	Sri Lanka	<i>Sogatella furcifera</i>	The extrusion of a puparium containing a male pupa or the opening of an adult female on the abdomen of the host is conspicuous as an external symptom of its parasitism. 40% parasitism rates reached – but was not persistent and failed to control the population of planthoppers	Ôtake et al, 1976
	Fiji, India, Indonesia, Japan, Philippines, Solomon Is., Sri Lanka, Thailand	<i>Sogatella furcifera</i> <i>Nilaparvata lugens</i> <i>Nephotettix virescens</i>		Greathead, 1982
	Malaysia	<i>Sogatella furcifera</i> <i>Nilaparvata lugens</i>	Claims <i>S. furcifera</i> and <i>N. lugens</i> were the two main planthoppers causing much damage in Malaysia. Suggests little was known about parasitoids of planthoppers at that time	Ooi, 1982

TABLE 3. ORDER: DIPTERA

Species name	Distribution Country/ region	Hosts	Notes (parasitism rates, alternative hosts, time of year, other information)	Reference (publication in English unless indicated otherwise)
Family: Pipunculidae				
Nymphal-adult parasites (Chandra, 1980)				
Favoured dryland environments (Chandra, 1980)				
No effective pipunculid parasitoid of <i>N. lugens</i> (Greathead, 1982)				
<i>Dorylas</i> sp.	Sri Lanka (citing Santa et al, unpublished)	<i>Nilaparvata lugens</i>		Chiu, 1979
<i>Pipunculus javanensis</i> de Meijere	Taiwan (citing Chiui et al, unpublished)	<i>Nilaparvata lugens</i>	low parasitism rates (citing Chiu, 1974)	Chiu, 1979
		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
	Taiwan	<i>Nephotettix cincticeps</i>	Nymphs and adults parasite	Chu & Hirashima,

		<i>Nilaparvata lugens</i>		1981
<i>Pipunculus mutillatus</i>		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
	India	<i>Nilaparvata lugens</i>	Nymph and adult parasite	Rhandhawa et al, 2006
	Thailand	<i>Nephotettix nigropictus</i> <i>Nephotettix virescens</i>		Wongsiri et al, 1980
	Taiwan	<i>Nephotettix cincticeps</i>	Nymphs and adults parasite	Chu & Hirashima, 1981
<i>Pipunculus orientalis</i>		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
<i>Pipunculus roralis</i>		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
<i>Pipunculus</i> Sp.	Sri Lanka, Taiwan	<i>Nilaparvata lugens</i> <i>Nephotettix cincticeps</i> <i>Nephotettix virescens</i>		Greathead, 1982
	Taiwan	<i>Nephotettix cincticeps</i> <i>Nephotettix nigropictus</i> <i>Nephotettix virescens</i>	Nymphs and adults parasite	Chu & Hirashima, 1981
<i>Tomosvaryella epichalca</i> Perkins	Taiwan (citing Chiui et al, unpublished)	<i>Nilaparvata lugens</i>	low parasitism rates (citing Chiu, 1974)	Chiu, 1979
		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
	Taiwan	<i>Nephotettix cincticeps</i> <i>Nilaparvata lugens</i>	Nymphs and adults parasite	Chu & Hirashima, 1981
<i>Tomosvaryella oryzaetoral</i> Koizumi	India	<i>Nilaparvata lugens</i>	Nymph and adult parasite	Rhandhawa et al, 2006
	Taiwan (citing Chiui et al, unpublished)	<i>Nilaparvata lugens</i>	low parasitism rates (citing Chiu, 1974)	Chiu, 1979
		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
	Zhejiang, Jiangxi, Fujian, Hubei	Green leafhopper		(no citation for this addition)
	Thailand	<i>Nephotettix nigropictus</i> <i>Nephotettix virescens</i>		Wongsiri et al, 1980
	Taiwan	<i>Nephotettix cincticeps</i> <i>Nilaparvata lugens</i>	Nymphs and adults parasite	Chu & Hirashima, 1981

<i>Tomosvaryella subvirescens</i> Loew	Taiwan, Thailand (citing Chiui et al, unpublished; Yasumatsu et al 1975)	<i>Nilaparvata lugens</i>	Important parasite of planthoppers and leafhoppers in Thailand – but population lower than in temperate countries In Taiwan – low parasitism rates (citing Chiu, 1974)	Chiu, 1979
		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
	Thailand	<i>Nephotettix nigropictus</i> <i>Nephotettix virescens</i>		Wongsiri et al, 1980
	Taiwan	<i>Nephotettix cincticeps</i> <i>Nilaparvata lugens</i>	Nymphs and adults parasite	Chu & Hirashima, 1981
<i>Tomosvaryella sylvatica</i>		<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Cab International, 2005
	Taiwan	<i>Nilaparvata lugens</i>	Nymphs and adults parasite	Chu & Hirashima, 1981

TABLE 4. ORDER: HEMIPTERA

Species name	Distribution Country/ region	Hosts	Notes (parasitism rates, alternative hosts, time of year, other information)	Reference (publication in English unless indicated otherwise)
Family: Anthocoridae				
<i>Orius tantillus</i> Motschulsky		<i>Nilaparvata lugens</i>	Nymph and adult parasite – this is listed in this reference as a parasitoid but other sources suggest it is a predator	CAB International, 2005

Comment [CSU7]: This species was listed by CABI as a parasitoid but other sources suggest it is a predator

Shankar & Baskaran (1988), in India, also report they frequently observed a mite belonging to the family Trombidiidae as a nymph and adult parasitoid of *N. lugens* .

Table 5. Parasitoids of *Cnaphalocrocis medinalis* (Guenée).

Species name	Family	Distribution Country/ region	Notes (parasitism rates, alternative hosts, time of year, other information)	Reference (publication in English unless indicated otherwise)
HYMENOPTERA				
<i>Apanteles anagleit</i>	Braconidae	India	Larval parasite	Randhawa et al, 2006
<i>Apanteles angustibasis</i>	Braconidae	India	Larval parasite	Randhawa et al, 2006
<i>Apanteles opacus</i> Ashmead	Braconidae	Peninsular Malaysia	Abundant parasitoid	van Vreden & Ahmadzabidi, 1986
<i>Apanteles cypris</i>	Braconidae	Taiwan	Host stage – larva Present in wet and dry seasons	Chou, 1981
	Braconidae	Peninsular Malaysia	Common parasitoid	van Vreden & Ahmadzabidi, 1986
	Braconidae	India	Larval parasite	Randhawa et al, 2006
<i>Apanteles opacus</i>	Braconidae	India	Larval parasite	Randhawa et al, 2006
<i>Apanteles syleptae</i>	Braconidae	India	Larval parasite	Randhawa et al, 2006
<i>Apanteles</i> sp.	Braconidae	Madurai district, India	Larval parasitoid	Rani et al, 2007
<i>Apanteles</i> sp. ater group	Braconidae	India	Larval parasitoid	Pati & Mathur, 1982
<i>Bracon</i> sp.	Braconidae	Philippines	Not important larval parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Bracon gelechiae</i>	Braconidae	India	Larval parasitoid	Randhawa et al, 2006

<i>Bracon hebetor</i>	Braconidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Bracon ricinicola</i>	Braconidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Cardiochiles philippinensis</i> Ashmead ²	Braconidae	Laguna, Philippines	Larval parasitoid - Present in wet and dry seasons	de Kraker et al, 1999
			Parasitism peaked at 28°C	Runjie et al, 1996 (abstract only)
			Common in dryland and wetland rice environments	http://www.knowledgebank.irri.org/Beneficials/Scientific_name/Cardiochiles_philippinensis_Ashmead_64.htm
		Philippines	Larval parasitoid	Ooi and Shepard, 1994
		Madurai district, India	Larval parasitoid	Rani et al, 2007
		Philippines	Very important larval parasitoid based on field collection and rearings	Barrion et al, 1991
		Tropical Asia	Larval parasitoid	Reissig et al, 1986
			Enters folded leaves and lays a single egg on leaffolder larva	Shepard et al, 2000
	India	Larval parasitoid	Randhawa et al, 2006	
<i>Cardiochiles laevifossa</i>	Braconidae	Taiwan	Host stage – larva Present in wet and dry seasons	Chou, 1981
<i>Cardiochiles</i> sp.	Braconidae	Peninsular Malaysia	Less common parasitoid	van Vreden & Ahmadzabidi, 1986
<i>Chelonus munakatae</i> Munakata	Braconidae	Philippines	Not important larval parasitoid based on field collection and rearings	Barrion et al, 1991
		Tropical Asia	Larval parasitoid	Reissig et al, 1986
	Braconidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Cotesia agilis</i> Ashmead	Braconidae		Hosts: hesperiid and rice leaffolder larvae (not sure if this means <i>C. medinalis</i> or not)	Barrion & Litsinger, 1994 p.209
<i>Cotesia cypris</i> Nixon	Braconidae	Tropical Asia	Larval parasitoid	Reissig et al, 1986
<i>Cotesia flavipes</i>	Braconidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Cotesia opacus</i> Ashmead	Braconidae		Alternative host: <i>Herpetogramma stultalis</i>	Barrion & Litsinger, 1994 p.207

<i>Cotesia ruficrus</i>	Braconidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Cotesia angustibasis</i> Gahan ²	Braconidae		Alternative hosts: <i>Marasmia</i> spp.	Barrion & Litsinger, 1994 p.207
		Philippines	Larval parasitoid	Ooi and Shepard, 1994
		Tropical Asia	Larval parasitoid	Reissig et al, 1986
		Peninsular Malaysia	Common parasitoid	van Vreden & Ahmadzabidi, 1986
		India	Larval parasitoid	Pati & Mathur, 1982
			Lays more than 10 eggs inside each leaffolder larval host. The hatched wasp larvae feed on the internal tissues of the host larva, eventually killing it. When ready to pupate, they leave the dead host and spin white cocoons nearby.	Shepard et al, 2000
<i>Cotesia</i> spp.	Braconidae	Philippines	Host stage – larva Present in wet and dry seasons	de Kraker et al, 1999
		Philippines	Larval parasitoid	de Kraker, 1996
		Philippines		Barrion et al, 1991

<i>Habrobacon</i> sp.	Braconidae	India	Larval parasitoid	Pati & Mathur, 1982
<i>Kriechbaumerella</i> sp.	Braconidae	India	Pupal parasitoid	Pati & Mathur, 1982
<i>Macrocentrus philippinensis</i> Ashmead ²	Braconidae	India	Extracts of <i>Saccharum officinarum</i> (sugarcane), <i>Cajanus cajan</i> (pigeonpea), <i>Oryza sativa</i> (rice), <i>Ricinus communis</i> (castor oil plant) and <i>Vigna sinensis</i> (field pea) affects parasitism and emergence (I do not know what host was used)	Shankarganesh & Khan, 2006 (abstract only)
			Alternative hosts: <i>Marasmia</i> spp.	Barrion & Litsinger, 1994 p.213
			Larval parasitoid	Ooi and Shepard, 1994
		Laguna, Philippines	Larval parasitoid – Present in wet and dry seasons	de Kraker et al, 1999
		India	2.0% parasitism rate	Rani et al, 2007
		Philippines	Very important larval parasitoid based on field collection and rearings	Barrion et al, 1991
		Tropical Asia	Larval parasitoid	Reissig et al, 1986
		Laid single egg on host larva and hatched as a single parasitoid larva	Shepard et al, 2000	

<i>Meteorus bacoorensis</i>	Braconidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Opius</i> sp.	Braconidae	Philippines	Not important larval parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Orgilus ashmeadii</i> Viereck	Braconidae		Alternative hosts: <i>Marasmia</i> spp.	Barrion & Litsinger, 1994 p.205
<i>Orgilus</i> sp.	Braconidae	Philippines	Moderately important larval parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Tropobracon schoenobii</i> Viereck	Braconidae	Philippines	Moderately important larval parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Goniozus indicus</i>	Bethylidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Goniozus</i> nr. <i>triangulifer</i> Kieffer ²	Bethylidae		Alternative hosts: <i>Marasmia</i> leaffolders	Barrion & Litsinger, 1994 p.184
		Philippines	Larval parasitoid	Ooi and Shepard, 1994
		Philippines	Very important larval parasitoid based on field collection and rearings	Barrion et al, 1991
			The wasp enters folded leaf and paralyzes the host larva before laying 3-8 eggs outside its body. The early-stage parasitoid larvae feed externally on leaffolder larva and then kill their host .	Shepard et al, 2000
<i>Goniozus triangulifer</i> Kieffer	Bethylidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Goniozus triangulus</i>	Bethylidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Goniozus</i> sp.	Bethylidae	Madurai district, India	Larval parasitoid 4.9% parasitism rate	Rani et al, 2007
		Philippines	Larval parasitoid	de Kraker, 1976
<i>Brachymeria excarinata</i> Gahan	Chalcididae	Peninsular Malaysia	Less common parasitoid	van vreden & Ahmadzabidi, 1986
		Tropical Asia	Larval parasitoid	Reissig et al, 1986
			Parasitizes older leaffolder larva	Shepard et al, 2000
		India	Larval /pupal parasitoid	Randhawa et al, 2006
<i>Brachymeria lasus</i> (Walker)	Chalcididae		Parasitizes older leaffolder larva	Shepard et al, 2000
		India	1.5% parasitism on pupal host	Bharati & Kushwaha, 1988
		India	Pupal parasitoid	Randhawa et al, 2006
<i>Brachymeria tacardiae</i>	Chalcididae	India	Pupal parasitoid	Randhawa et al, 2006

<i>Brachymeria sp. cf. tarsalis</i> (Motschulsky)	Chalcididae	Peninsular Malaysia	Less common parasitoid	van Vreden & Ahmadzabidi, 1986
<i>Brachymeria sp./spp.</i>	Chalcididae	Madurai district, India	Larval parasitoid 6.5% parasitism rate	Rani et al, 2007
			Parasitizes older leaffolder larva	Shepard et al, 2000
		Philippines	Very important larval-pupal parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Trachichospilus pupivora</i>	Chalcididae	India	Pupal parasitoid	Randhawa et al, 2006
<i>Elasmus brevicornis</i>	Elasmidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Elasmus claripennis</i>	Elasmidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Elasmus philippinensis</i> Ashmead	Elasmidae	Peninsular Malaysia	Larval parasitoid	van Vreden & Ahmadzabidi, 1986
		India	Larval parasitoid	Randhawa et al, 2006
<i>Elasmus sp./spp.</i>	Elasmidae	Philippines	Larval parasitoid	de Kraker, 1996
		Tropical Asia	Larval parasitoid	Reissig et al, 1986
			One or two eggs are laid in each young or old larva. They are highly aggressive and will kill other parasitoid larvae, which may be developing inside the leaffolder host.	Shepard et al, 2000
		Philippines	Not important larval parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Copidosoma sp.</i>	Encyrtidae	India	Egg parasite	Encyrtidae
<i>Copidosomopsis nacoleiae</i> Eady ¹	Encyrtidae		Alternative hosts: <i>Marasmia</i> spp., <i>Hedylepta indicata</i>	Barrion & Litsinger, 1994 p.266
		Philippines	Egg-larval parasitoid	Ooi and Shepard, 1994
			Larval parasitoid – present in wet and dry seasons	de Kraker et al, 1999
		Madurai district, India	Larval parasitoid 3.1% parasitism rate	Rani et al, 2007
		Philippines	Egg-larval parasitoid	de Kraker, 1996
	Philippines	Very important egg-larval parasitoid based on field collection and rearings	Barrion et al, 1991	

			200-300 wasps are produced from a few host eggs and hundreds of wasp pupae can be seen through the skin of the host larva	Shepard et al, 2000
		Tropical Asia	Egg-larval parasitoid	Reissig et al, 1986
		India	Egg parasite	Randhawa et al, 2006
<i>Tetrastichus ayyari</i> Rohwer ²	Eulophidae	Philippines	Pupal parasitoid	Ooi and Shepard, 1994
<i>Tetrastichus howardi</i> Olliff (= <i>ayyari</i> Rohwer)	Eulophidae	Philippines	Not important pupal parasitoid based on field collection and rearings	Barrion et al, 1991
		India	Pupal parasitoid	Randhawa et al, 2006
<i>Tetrastichus israelensis</i>	Eulophidae	India	Pupal parasitoid	Randhawa et al, 2006
<i>Tetrastichus schoenobii</i> Ferriere	Eulophidae	Philippines	Not important larval-pupal parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Tetrastichus</i> sp.	Eulophidae	Philippines	Pupal parasitoid	de Kraker, 1996
<i>Stenomesus</i> sp.	Eulophidae	Philippines	Not important larval parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Aphanogmus fijiensis</i>	Ichneumonidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Barylypa apicala</i>	Ichneumonidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Charops brachypterum</i> Cameron	Ichneumonidae		Alternative hosts: <i>Marasmia</i> spp., <i>S.innotata</i>	Barrion & Litsinger, 1994 p.227
		Philippines	Not important larval parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Charops nigrita</i> Gupta & Maheswary	Ichneumonidae	Philippines	Not important larval parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Diatora lissonata</i>	Ichneumonidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Eribborus argenteopilosus</i>	Ichneumonidae	India	Larval/pupal parasitoid	Randhawa et al, 2006
<i>Eriborus sinicus</i> (Holmgren)	Ichneumonidae	India	Larval/pupal parasitoid	Randhawa et al, 2006
		Philippines	Not important larval parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Ischnojoppa luteator</i> (Fabricius)	Ichneumonidae	Philippines	Very important larval parasitoid based on field collection and rearings	Barrion et al, 1991
		India	Larval/pupal parasitoid	Randhawa et al, 2006

<i>Itopectis narangae</i> (Ashmead)	Ichneumonidae	Philippines	Very important larval parasitoid based on field collection and rearings	Barrion et al, 1991
			A solitary hunter, can locate larvae inside stems	Shepard et al, 2000
<i>Leptobatopsis indica</i>	Ichneumonidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Stictopisthus</i> sp.	Ichneumonidae	Philippines	Not important larval parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Temelucha basimacula</i>	Ichneumonidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Temelucha biguttula</i> Manakata	Ichneumonidae		Alternative hosts: <i>C. suppressalis</i> , <i>Naranga diffusa</i> , <i>Bradina admixtalis</i>	Barrion & Litsinger, 1994 p.225
		Guangdong Province, China	A common parasitoid species found in that area	Ying, 1982
		India	Larval parasitoid	Randhawa et al, 2006
<i>Temelucha philippinensis</i> Ashmead ²	Ichneumonidae		Alternative hosts: <i>C. suppressalis</i> , <i>Marasmia</i> spp.	Barrion & Litsinger, 1994 p.225
		Philippines	Larval parasitoid	Ooi and Shepard, 1994
			Larval parasitoid – present in wet and dry seasons	de Kraker et al, 1999
		Guangdong Province, China	A common parasitoid species found in that area	Ying, 1982
		Tropical Asia	Larval parasitoid	Reissig et al, 1986
			Hunts leaffolder larva during the day	Shepard et al, 2000
		Philippines	Very important larval parasitoid based on field collection and rearings	Barrion et al, 1991
		India	Larval parasitoid	Randhawa et al, 2006
<i>Temelucha stangli</i> Ashmead	Ichneumonidae		Alternative hosts: <i>Scirpophaga</i> spp., <i>Marasmia</i> spp.	Barrion & Litsinger, 1994 p.225
		Guangdong Province, China	A common parasitoid species found in that area	Ying, 1982
		Philippines	Very important larval parasitoid based on field collection and rearings	Barrion et al, 1991
		Tropical Asia	Larval parasitoid	Reissig et al, 1986
		India	Larval parasitoid	Randhawa et al, 2006
<i>Trathala flavo-orbitalis</i> (Cameron)	Ichneumonidae	Philippines	Moderately important larval parasitoid based on field collection and rearings	Barrion et al, 1991
		India	Larval/pupal parasitoid	Randhawa et al, 2006
<i>Trichomma</i>	Ichneumonidae		Alternative hosts: <i>Marasmia</i> spp., <i>Ostrinia furnacalis</i>	Barrion & Litsinger, 1994

<i>cnaphalocrosis</i> Uchida ²				p.225
		Philippines	Larval parasitoid	Ooi and Shepard, 1994
			Larval/pupal parasitoid – present in wet and dry seasons	de Kraker et al, 1999
		Madurai district, India	Larval parasitoid 10.4% parasitism rate	Rani et al, 2007
		Tropical Asia	Larval parasitoid	Reissig et al, 1986
		Philippines	Very important larval parasitoid based on field collection and rearings	Barrion et al, 1991
		Tropical Asia	Larval parasitoid	Reissig et al, 1986
<i>Xanthopimpla flavineata</i> Cameron ²	Ichneumonidae		Alternative hosts: <i>C. supressalis</i> , <i>S. inferens</i> , <i>P. Mathias</i> , <i>Telicota augias</i>	Barrion & Litsinger, 1994 p.215
		Philippines	Pupal parasitoid	Ooi and Shepard, 1994
		Madurai district, India	Pupal parasitoid 0.7% parasitism rate	Rani et al, 2007
		India	Pupal parasitoid with 20-23.3% parasitism rate	Bharati & Kushwaha, 1988
		India	Pupal parasitoid	Pati & Mathur, 1982
		Philippines	Not important pupal parasitoid based on field collection and rearings	Barrion et al, 1991
		India	Pupal parasitoid	Randhawa et al, 2006
<i>Trichomalopsis apanteloctena</i> (Crawford)	Pteromalidae	Philippines	Not important pupal parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Telenomus digmus</i>	Scellionidae	India	Egg parasite	Randhawa et al, 2006
<i>Trichogramma chilonis</i> Ishii,	Trichogrammati dae	Pakistan	Neem seed kernel extract and <i>Bacillus thurigiensis</i> may also be used to enhance its efficacy of the parasitoid	Sagheer et al, 2008
		India	Alternative hosts: <i>Helicoverpa armigera</i> , <i>Spodoptera litura</i> and <i>Papilio demoleus</i> (out of these preferred <i>H. armigera</i>)	Budhwant et al, 2008 (abstract only)
		India	Extracts of <i>Saccharrum offinarum</i> (sugarcane), <i>Cajanus cajan</i> (pigeonpea), <i>Oryza sativa</i> (rice), <i>Ricinus communis</i> (castor oil plant) and <i>Vigna sinensis</i> (field pea) affects parasitism and emergence (I do not know what host was used)	Shankarganesh & Khan, 2006 (abstract only)
		India	Indundative release provided effective protection against <i>Scirpophaga incertulas</i> and <i>C. Medinalis</i> with 100,000/ha better than lower doses	Kumar & Khan, 2005 (abstract only)
<i>Trichogramma</i>	Trichogrammati	India	Indundative release provided effective protection against	Kumar & Khan, 2005

<i>japonicum</i> Ashmead ¹	dae		<i>Scirpophaga incertulas</i> and <i>C. Medinalis</i> with 100,000/ha better than lower doses	(abstract only)
		Philippines	Egg parasitoid Alternative host: <i>Chilo suppressalis</i> (citing Sweezy, 1931)	Ooi and Shepard, 1994
		Japan, Hawaii	Alternative host: <i>Chilo simplex</i>	Sweezy, 1931
		Guangdong Province, China	A common parasitoid species found in that area	Ying, 1982
		Philippines	Egg parasitoid	de Kraker, 1996
		Philippines	Moderately important egg parasitoid based on field collection and rearings	Barrion et al 1991
<i>Trichogramma poliae</i>		India	Extracts of <i>Saccharum officinarum</i> (sugarcane), <i>Cajanus cajan</i> (pigeonpea), <i>Oryza sativa</i> (rice), <i>Ricinus communis</i> (castor oil plant) and <i>Vigna sinensis</i> (field pea) affects parasitism and emergence (I do not know what host was used)	Shankarganesh & Khan, 2006 (abstract only)
<i>Trichogramma</i> sp./spp.	Trichogrammati dae	India	Used for inundative biocontrol – listed as an important natural enemy of leaf folder	Pasalu et al, 2004
		Madurai district, India	Egg parasite 9.3% parasitism rate	Rani et al, 2007
		Tropical Asia	Egg parasitoid	Reissig et al, 1986
		Philippines	Very important egg parasitoid based on field collection and rearings	Barrion et al, 1991
		Philippines	Egg stage mortality in the field averaged about 60% due to parasitism	de Kraker, 1996
DIPTERA				
<i>Megaselia scalaris</i>	Phoridae	India	Larval parasitoid	Randhawa et al, 2006
<i>Megaselia</i> sp.	Phoridae	Philippines	Moderately important larval parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Argyrophylox fransseni</i> Baranov	Tachinidae	Peninsular Malaysia	Reared as parasitoid from <i>C. medinalis</i>	van Vreden & Ahmadzabidi, 1986
<i>Argyrophylox nigrotibialis</i> (Baranov)	Tachinidae	Philippines	Moderately important larval parasitoid based on field collection and rearings	Barrion et al, 1991
<i>Chaetexorista javana</i>	Tachinidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Nemorilla floralis</i>	Tachinidae	India	Larval parasitoid	Randhawa et al, 2006
<i>Zygothria ciliata</i>	Tachinidae	Philippines	Moderately important larval parasitoid based on field	Barrion et al, 1991

Comment [CSU8]: Not sure if this parasitoid affects this pest

(Wulp)		collection and rearings	
--------	--	-------------------------	--

¹ Important according to Ooi and Shepard (1994)

² common according to Ooi and Shepard (1994)

References

(websites not listed – follow hyperlinks for more information)

- Baquero E & Jordana R (1999) Species of *Anagrus* Haliday, 1833 (Hymenoptera, Chalcidoidea, Mymaridae) in Navarra (Spain), *Miscel-lània Zoològica*, **22** (2): 39-50
- Barrion AT, Pantua PC, Bandong JP, dela Cruz CG, Raymundo FA, Lumaban MD, Apostol RF & Litsinger JA (1981) Food web of the rice brown planthopper in the Philippines, *International Rice Research Notes*, **6** (1): 13-15
- Barrion AT & Litsinger JA, Medina EB, Aguda RM, Bandong JP, Pantua Jr. PC, Viajante VD, dela Cruz CG, Vega CR, Soriano Jr. JS, Camanag EE, Saxena RC, Tryon EH, Shepard BM (1991) The rice *Cnaphalocrocis* and *Marasmia* (Lepidoptera :Pyrallidae) leafroller complex in the Philippines : Taxonomy, bionomics and control, *Philippine Entomologist*, **8** (4): 987-1074
- Barrion AT & Litsinger JA (1994) Taxonomy of rice insect pests and their arthropod parasites and predators. In: EA Heinrichs (ed) *Biology and management of rice insects*, New York: John Wiley & Sons Inc, pp. 13-359
- Bharati LR & Kushwaha KS (1988) Parasitoids of leafroller (LF) pupae from Haryana, India, *International Rice Research Newsletter*, **13** (5):31
- Budhwant NP, Dadmal SM, Nemade PW & Patil MS (2008) Efficacy of *Trichogramma chilonis* Ishii against lepidopteron pests and age of host eggs, *Annals of Plant Protection Sciences*, **16** (1): (no page numbers)
- CAB International (2005) Crop Protection Compendium, 2005 Edition, Wallingford, UK: CAB International. www.cabicompendium.org/cpc Accessed: 6/08/2009
- Chandra G (1980) Taxonomy and bionomics of the insect parasites of rice leafhoppers and planthoppers in the Philippines and their importance in natural biological control, *Philippine Entomologist*, **4** (3): 119-139
- Chandra G & Dyck VA. (1987) New techniques of rearing *Anagrus flaveolus* (Hym.:Mymaridae), an egg parasitoid of the brown planthopper, *BioControl*, **33** (2): 239-243
- Chantarasa-ard S (1984) Preliminary study on the overwintering of *Anagrus incarnates* Haliday (Hymenoptera: Mymaridae), an egg parasitoid of the rice planthoppers, *Esakia*, **22**: 15-162
- Chantarasa-ard S & Hirashima Y (1984) Host range and host suitability of *Anagrus incarnatus* Haliday (Hymenoptera: Mymaridae), an egg parasitoid of Delphacid planthoppers, *Applied Entomology and Zoology*, **19** (4): 491-497
- Chantarasa-ard SJ, Hirashima Y & Miura T (1984) Effects of temperature and food on the development and reproduction of *Anagrus incarnatus* Haliday (Hymenoptera : Mymaridae): an egg parasitoid of the rice planthoppers, *Esakia*, **22**: 145-158
- Charleston DS, Rami K, van Rensburg NJ, Barnes BN, Hattingh V, Conlong DE, Visser D & Prinsloo GJ (2003) Intergrated pest management in South Africa. In: KM Maredia, D Dakouo & D Mota-Sanchez (eds), *Integrated Pest Management in the Global Arena*, CABI Publishing: Wallingford, pp. 169-195
- Chen B-H & Yu, JZ (1989) *Anagrus incarnates* Haliday, a new record from eggs of Brown Planthopper in Taiwan, *Journal of Agricultural Research China*, **38** (4): 458-462
- Chen M, Ye GY, Hu C & Datta SK (2003) Effect of transgenic Bt rice on dispersal of planthoppers and leafhoppers as well as their egg parasitic wasps, *Journal of Zhejiang University (Agriculture and Life Sciences)*, **29** (1): 29-33
- Chen Y & Hirashima Y (1981) Survey of Taiwanese literature on the natural enemies of rice leafhoppers and planthoppers, *Esakia*, **16**: 33-37

- Chen YX (1989) Investigation of dryinids of rice planthoppers in Sinan County, Guizhou, China, *Insect Knowledge*, **26** (2): 77-79
- Chiappini E, Soressi L, Fogher C & Zanirato M (1999) Genetic identity and relationship between four *Anagrus species* (Hymenoptera: Mymaridae) using RAPD analysis, *European Journal of Entomology*, **96**: 393-400
- Chiappini E & Lin N-Q (1998) *Anagrus* (Hymenoptera: Myrmaridae) of China, with descriptions of nine new species, *Annals of the Entomological Society of America*, **91** (5): 549-571
- Chiu C-S (1979) Biological control of the brown planthopper. In: International Rice Research Institute, *Brown planthopper: threat to rice production in Asia*, Manila: International Rice Research Institute, pp.335-355
- Chou L-Y (1981) A preliminary list of Braconidae (Hymenoptera) of Taiwan, *Journal of Agricultural Research China*, **20** (1): 71-88
- Chua TH, Dyck VA & Peña NB (1984) Functional response and searching efficiency in *Pseudogonatopus flavifemur* Esaki and Hash. (Hymenoptera: Dryinidae), a parasite of rice planthoppers, *Researches on Population Ecology*, **26**: 74-83
- Claridge MF (1996) *Biological control of brown planthopper in Indonesia*. In: biotec@goliat.ugr.es.
- Claridge MF, Morgan JC, Steenkiste AE, Iman M & Damayanti D (1999) Seasonal patterns of egg parasitism and natural biological control of rice brown planthopper in Indonesia, *Agricultural and Forest Entomology*, **1** (4):297-304
- Dayanan GJ & Esteban EP (1996) Natural enemies of brown planthopper *Nilaparvata lugens* Stal in lowland rice at USM, 27th Anniversary and Annual Scientific Meeting of the Pest Management Council of the Philippines, Inc., Davao City, Philippines, 7-10 May 1996
- de Kraker J (1996) *Impact of natural enemies on rice leafhopper*, Department of Entomology, Wageningen Agricultural University. 32 p.
- de Kraker J (1996) *The potential of natural enemies to suppress rice leafhopper populations*, Department of Entomology, Wageningen Agricultural University. 257 p.
- de Kraker J, van Huis A, Heong KL, van Lenteren JC & Rabbinge R (1999) Population dynamics of rice leafhoppers (Lepidoptera: Pyralidae) and their natural enemies in irrigated rice in the Philippines, *Bulletin of Entomological Research*, **89**: 411-421
- Dick J & Thomson GM (1969) The effect of a systemic insecticide on the spread of mosaic and on certain sugarcane insects, *Proceedings of the South African Technologists' Association*, **June**, 72-74
- Fowler SV, Claridge MF, Morgan JC, Peries IDR & Nugaliyadde L (1991) Egg mortality of the brown planthopper, *Nilaparvata lugens* (Homoptera: Delphacidae) and green leafhoppers, *Nephotettix* spp. (Homoptera: Cicadellidae), on rice in Sri Lanka, *Bulletin of Entomological Research*, **81**: 161-167
- Greathead DJ (1982) Natural enemies of *Nilaparvata lugens* and other leaf- and plant-hoppers in tropical agroecosystems and their impact on pest populations. In: WJ Knight, NC Pant, TS Robertson & MR Wilson (eds), Proceedings of the 1st International Workshop on Biotaxonomy, *Classification and Biology of Leafhoppers and Planthoppers (Auchenorrhyncha) of Economic Importance*, London, 4-7 October, Commonwealth Institute of Entomology: London
- Guerrieri E & Viggiani G (2005) A review of the encyrtid (hymenoptera: Chalcidoidea) parasitoids of Dryinidae (Hymenoptera: Chrysidoidea) with descriptions of a new species of *Cheiloneurus*, *Systematics and Biodiversity*, **2** (3): 305-317
- Gupta A & Poorani J (2008) New distribution and host records of Chalcidoidea (Insecta: Hymenoptera) from various parts of India, *Check List*, **4** (4): 410-414
- Gupta M & Pawar AD (1989) Biological control of rice leafhoppers and planthoppers in Andhra Pradesh, *Plant Protection Bulletin (Faridabad)*, **41** (1-2): 6-11
- Hachiya K (1995) Egg parasitoid of small brown planthopper, *Laodelphax striatellus* Fallen (Homoptera, Delphacidae) in Hokkaido, Japan, *Japanese Journal of Applied Entomology and Zoology*, **39** (1): 89-90
- Haug SS (1994) The construction of the life table of whitebacked planthopper, *Journal of South China Agricultural University*, **15** (3): 45-49 (in Chinese – English abstract)
- Heong KL, Aquino GB & Barrion AT (1992) Population dynamics of plant- and leafhoppers and their natural enemies in rice ecosystems in the Philippines, *Crop Protection*, **11**: 371-379
- Hiroshima Y (1978) Strepsipterous parasites of homoptera injurious to the rice plant in Sarawak, Borneo, with description of a new species, *Esakia*, **11**: 53-58

Comment [CSU9]: Does not distinguish between these two references

- Hiroshima Y (1979) Field studies on the biological control of leafhoppers and planthoppers (Hemiptera: Homoptera) injurious to rice plants in South-East Asia. Progress report for the year 1977, *Esakia*, **13**: 1-20
- Hu JS (1992) Observations on the spatial habits of two species of *Anagrus* parasites, *Insect Knowledge*, **29** (2): 107-109
- Ito E & Yamanda YY (2007) Imperfect preference for singly parasitized hosts over doubly parasitized hosts in the semisolitary parasitoid *Echthrodelphax fairchildii*: implications for profitable self-superparasitism, *Entomologia Experimentalis et Applicata*, **123**: 207-215
- Kitamura K (1982) Comparative studies on the biology of dryinid wasps in Japan. 1. Preliminary report on the predacious and parasitic efficiency of *Halogonaptus atratus* Esaki et Hashimoto (Hymenoptera: Dryinidae), *Bulletin of the Faculty of Agriculture, Shimane University*, **16**: 172-176
- Kitamura K (1987) Seasonal changes in percentage parasitism of the parasitoids of leaf- and planthoppers in Shimane pref. (Homoptera: Auchenorrhyncha), *Bulletin of the Faculty of Agriculture, Shimane University*, **21**: 155-157
- Kitamura K (1989) Comparative studies on the biology of dryinid wasps in Japan. 10. Development of *Psuedogonatopus fulgori* (Hymenoptera, Dryinidae), *Shimane Daigaku Nogakuba Kenkyu – Bulletin of the Faculty of Agriculture, Shimane University*, **23**: 60-63
- Kitamura K & Iwami J (1998) Comparative studies on the biology of dryinid wasps in Japan. 14. Sex allocation and size of *Haplogonatopus apicalis* (Hymenoptera: Dryinidae), in relation to instars of host nymphs and host sex, *Japanese Journal of Entomology* (New Series), **1** (1): 1-8
- Kumar S & Khan MA (2005) Bio-efficacy of *Trichogramma* spp. against yellow stem borer and leaf folder in rice ecosystem, *Annals of Plant Protection Sciences*, **13** (1): (no page numbers)
- Li BC & He JX (1991) Investigation of population dynamics of three species of Mymaridae parasitizing the eggs of planthoppers and their utilization and protection, *Natural Enemies of Insects*, **13** (4): 156-161
- Li, L-Y (1982) Integrated rice insect pest control in the Guangdong Province of China, *Entomophaga (Biocontrol)*, **27** (Special Issue): 81-88
- Liljeström GG & Virla EG (2004) Density-dependent parasitism of *Delphacodes kusheli* eggs by *Anagrus flaveolus*: influence of egg patchiness and density, *Biocontrol Science and Technology*, **14** (2): 107-115
- Lo SN & Zhou WX (1980) Investigations on the bionomics and utilization of mymarid egg-parasites of rice planthoppers, *Journal of Fujian Agricultural College*, **2**: 44-60
- Lou Y-G & Cheng, J-A (2001) Host-recognition kairomone from *Sogatella furcifera* for the parasitoid *Anagrus nilaparvatae*, *Entomologia Experimentalis et Applicata*, **101**: 59-67
- Lou Y-G, Du M-H, Turlings TCJ, Cheng J-A & Shen, W-F (2005a) Exogenous application of jasmonic acid induces volatile emissions in rice and enhances parasitism of *Nilaparvata lugens* eggs by the parasitoid *Anagrus nilaparvatae*, *Journal of Chemical Ecology*, **31** (9): 1985-2002
- Lou Y-G, Ma B, Cheng J-A (2005b) Attraction of the parasitoid *Anagrus nilaparvatae* to rice volatiles induced by the rice brown planthopper *Nilaparvata lugens*, *Journal of Chemical Ecology*, **31** (10): 2357-2372
- Lou X & Zhou W (1986) Studies on the relationships of population fluctuation between rice planthoppers and natural enemies and natural control effects, *Natural Enemies of Insects*, **8** (2): 72-79
- Lu Y, Wang X, Lou Y & Cheng J (2006) Role of ethylene signalling in the production of rice volatiles induced by the rice brown planthopper *Nilaparvata lugens*, *Chinese Science Bulletin*, **51** (20): 2457-2465
- Mao RQ, Zhang WQ, Zhang GR & Gu DX (2002a) An investigation on dynamics of egg-parasitoid community on planthopper in rice area in South China, *Acta Entomologica Sinica*, **45** (1): 96-101
- Mao RQ, Gu DX, Zhang GR & Zhnag WQ (2002b) A preliminary investigation on structure and dynamics of egg parasitoid community on the brown planthopper in rice field, *Acta Entomologica Sinica*, **45** (3): 408-412
- Manjunath TM, Rai PS & Gowda G (1978) Parasites and predators of *Nilaparvata lugens* in India, *PANS*, **24** (3): 265-269

- Miura T, Hirashima Y & Wongsiri T (1979) Egg and nymphal parasites of rice leafhoppers and planthoppers. A result of field studies in Thailand in 1977, *Esakia*, **13**: 21-44
- Miura T, Hirashima YH, Ctib MT & Yau IC (1981) Egg and nymphal parasites of rice leafhoppers and planthoppers, a result of field studies in Taiwan in 1979 (Part 1), *Esakia*, **16**: 39-50
- Nalini R (2005) Interaction among resistant rice genotypes, whitebacked planthopper *Sogatella furcifera* (Horvath), and egg parasitoid *Anagrus nr. flaveolus*, *International Rice Research Notes*, **30** (1):29-30
- Olmi M (1991-92) Contribution to the knowledge of the Gonatopodinae (Hymenoptera Dryinidae), *Bollettino dell'Instituto di Entomologia "Guido Grandi" della Universita degli Studi di Bologna.*, **46**: 109-122
- Ooi PAC (1982) Attempts at forecasting rice planthopper populations in Malaysia, *Entomophaga* 27 (Special Issue): 89-98
- Ooi PAC & Shepard BM (1994) Predators and parasitoids of rice insect pests. In: EA Heinrichs (ed) *Biology and management of rice insects*, New York: John Wiley & Sons Inc, pp. 585-612
- Ôtake A (1970) Studies on the egg parasites of the smaller brown planthopper, *Laodelphax striatellus* (Fallen) (Hemiptera: Delphacidea). IV. Seasonal trends in parasitic and dispersal activities, with special reference to *Anagrus nr. flaveolus* Waterhouse (Hymenoptera: Mymaridae), *Applied Entomology & Zoology*, **5** (2):95-104
- Ôtake A (1977) Natural enemies of the brown planthopper. In: Asia and Pacific Council: Food Fertilizer Technology Center (compilers) *The brown planthopper*, ASPAC: food and Fertilizer Technology Center: Taipei, Taiwan, pp. 42-56
- Ôtake A, Somasundaram PH & Abeykoon MB (1976) Studies on populations of *Sogatella furcifera* Horvath and *Nilaparvata lugens* Stål (Hemiptera: Delphacidae) and their parasites in Sri Lanka, *Applied Entomology & Zoology*, **11** (3): 284-294
- Pasalu IC, Mishra B, Krishnaiah NV & Katti G (2004) Integrated pest management in rice in India: status and prospects. In: PS Birtal & OP Sharma (eds), *Integrated pest management in Indian agriculture: Proceedings 11*, National Centre for Agricultural Economics and Policy Research: New Delhi and National Centre for Integrated Pest Management: New Delhi, pp.25-37
- Pati P & Mathur KC (1982) New records of parasitoids attacking rice leafhopper, *Cnaphalocrocis medinalis* Guenee in India. *Current China*. **51** (18): 904-905
- Peña N & Shepard M (1986) Seasonal innce of parasitism of brown planthoppers *Nilaparvata lugens* (Homoptera: Delphacidae), green leafhoppers, *Nephotettix* spp., and whitebacked planthoppers, *Sogatella furcifera* (Homoptera: Cicadellidae), *Environmental Entomology*, **15**: 263-267
- Randhawa GJ, Bhalla S, Chalam VC, Tyagi V, Verma DD & Hota M (2006) *Document on biology of rice (Oryza sativa L.) in India*, National Bureau of Plant Genetic Resources: New Delhi
- Rani WB, Amutha R, Muthulakshmi S, Indira K & Mareeswari (2007) Diversity of rice leaf folders and their natural enemies, *Research Journal of Agriculture and Biological Sciences*, **3** (5): 394-397
- Reissig WH, Heinrichs EA, Litsinger JA, Moody K, Fiedler L, Mew TW & Barrion AT (1986) *Illustrated guide to integrated pest management in rice in tropical Asia. Manila (Philippines): International Rice Research Institute. 411 p.*
- Rhandhawa GJ, Bhalla S, Chalam VC, Tyagi V, Verma DD & Hota M (2006) Document on biology of rice (*Oryza sativa* L.) in India, National Bureau of Plant Genetic Resources: New Delhi
- Runjie Z, Heong KL & Domingo IT (1996) Relationship between temperature and functional response in *Cardiochiles philippinensis* (Hymenoptera: Braconidae), a larval parasitoid of *Cnaphalocrocis medinalis* (Lepidoptera: Pyralidae), *Environmental Entomology*, **25** (6): 1321-1324
- Sagheer M, Ashfaq M, Mansoor-ul-Hasan & Rana SA (2008) Integration of some biopesticides and *Trichogramma chilonis* for the sustainable management of rice leaf folder, *Cnaphalocrocis medinalis* (Guenee) (Lepidoptera: Pyralidae), *Pakistani Journal of Agricultural Science*, **45** (1): 69-74
- Sahad KA (1984) Biology of *Anagrus optabilis* (Perkins) (Hymenoptera: Mymaridae) an egg parasitoid of delphacid planthoppers, *Esakia*, **22**: 129-144

- Sahragard A, Jervis MA & Kidd NAC (1991) Influence of host availability on rates of oviposition and host-feeding, and on longevity in *Dicondylus indianus* Olmi (Hym., Dryinidae), a parasitoid of the Rice Brown Planthopper, *Nilaparvata lugens* Stål (Hem., Delphacidae), *Journal of Applied Entomology*, **112**: 153-162
- Shankar G & Baskaran P (1988) Impact of the presence of parasites on the population of resident endosymbiotes in brown planthopper, *Nilaparvata lugens* (Stål) (Delphacidae: Homoptera), *Current Science*, **57** (4):212-214
- Shankar G & Baskaran P (1992) Regulation of yeast-like endosymbiotes in the rice brown planthopper *Nilaparvata lugens* Stal (O: Homoptera, F: Delphacidae), *Symbiosis (Rehovot)*, **14** (1-3): 161-173
- Shankarganesh K & Khan MA (2006) Bio-efficacy of plant extracts on parasitisation of *Trichogramma chilonis* Ishii, *T. japonicum* Ashmead and *T. poliae* Nagaraja, *Annals of Plant Protection Sciences*, **14** (2): (no page numbers)
- Shepard BM, Barrion AT & Litsinger JA (2000) *Friends of the rice farmer: helpful insects, spiders, and pathogens*. Manila (Philippines): International Rice Research Institute. 136 p
- Singh SP, Rao NS & Henneberry TJ (1993) Leafhoppers and their natural enemies, *Technical Bulletin – Project Directorate of Biological Control, ICAR*, **6**, 65pp.
- Subba Rao BR (1976) *Narayana*, gen. nov. from Burma and some synonyms (Hymenoptera: Mymaridae), *Oriental Insects*, **10** (1): 87-91
- Swezey OH (1925) Records of introduction of beneficial insects into the Hawaiian Islands, *The Hawaiian Planters' Record*, **29** (4): 369-376
- Swezey OH (1931) Some recent parasite introductions in Hawaii, *Journal of Economic Entomology*, **24**: 945-947
- Tao C & Ngoan N (1970) An ecological study of white-back planthopper, *Sogatella furcifera* Horvath in Vietnam, 1968. *Chinese Electronic Periodical Services*, **19** (4): 82-90
- Triapitsyn SV (2001) Review of the Australasian species of *Anagrus* (Hymenoptera: Mymaridae), *Belgian Journal of Entomology*, **3** (2): 267-289
- Triapitsyn SV & Beardsley JW (2000) A review of the Hawaiian species of *Anagrus* (Hymenoptera: Mymaridae), *Proceedings of the Hawaiian Entomological Society*, **34**: 23-48
- van Vreden G & Ahmadzabidi AL (1986) *Pests of rice and their natural enemies in Peninsular Malaysia*. Wageningen (Netherlands): Centre for Agricultural Publishing and Documentation (Pudoc). 230 p.
- Vungsilabutr P (1981) Relative composition of egg-parasite species of *Nilaparvata lugens*, *Sogatella furcifera*, *Nephotettix virescens* and *N. nigropictus* in paddy fields in Thailand, *Tropical Pest Management*, **27** (3): 313-317
- Vungsilabutr P, Miura T & Miura K (1996) Can *Paracetrotia andoi* Ishii (Hymenoptera, Trichogrammatidae) parasitize the egg of the brown planthopper? *Japanese Journal Of Entomology*, **64** (4): 876-878
- Watanabe T, Wada T, Mohd N, Noor, B & Salleh, N (1992) Parasitic activities of egg parasitoids on the rice planthoppers, *Nilaparvata lugens* (Stål) and *Sogatella furcifera* (Horváth) (Homoptera: Delphacidae), in the Muda Area of Peninsular Malaysia, *Applied Entomology and Zoology*, **27** (2): 205-211
- Wongsiri T, Wongsiri C, Navavichit S, Lewvanich & Yatsumatsu K (1980) Abundance of natural enemies of rice insect pests in Thailand, *Proceedings of the Symposium on Tropical Agricultural Research*, **6-7**: 131-149
- Xiang, C Ren, N, Wang X, Sumera A, Cheng J & Lou, Y (2008) Preference and performance of *Anagrus nilaparvatae* (Hymenoptera: Mymaridae): effect of infestation duration and density by *Nilaparvata lugens* (Homoptera: Delphacidae), *Environmental Entomology*, **37** (3): 748-754
- Yadav KP & Pawar AD (1989) New record of dryinid parasitoid of brown planthopper, *Nilaparvata lugens* Stal and whitebacked planthopper, *Sogatella furcifera* Horv, *Entomon*, **14**, 3-4: 369-370
- Yamada YY and Ikawa K (2003) Adaptive significance of facultative infanticide in the semi-solitary parasitoid *Ecthodolapha fairchildii*, *Ecological Entomology*, **28**: 613-621
- Yamada YY & Kawamura (1999) Sex identification of eggs of a dryinid parasitoid, *Halogonatopus atratus*, based on oviposition behaviour, *Entomologia Experimentalis et Applicata*, **93**: 321-324

- Yamada YY & Kitashiro S (2002) Infanticide in a dryinid parasitoid, *Journal of Insect Behaviour*, **15** (3): 415-427
- Yamada YY & Miyamoto K (1998) Payoff from self and conspecific superparasitism in a dryinid parasitoid, *Haplogonatopus atratus*, *Oikos*, **81** (2):209-216
- Yu XP (1996) Relationship of egg parasitoids of rice planthoppers between rice and non-rice habitats, *Chinese Rice Newsletter, CRRN*, **4** (3): 9-11
- Yu XP, Hu C & Heong KL (1996) Effects of non-rice habitats on the egg parasitoids of rice planthoppers, *Journal of Zhejiang Agricultural University*, 1996, **22** (2): 115-120
- Yu XP, Hu C & Heong KL (1998) Parasitization and preference characteristics of egg parasitoids from various habitats to homopterans, *Acta Entomologica Sinica*, **41** (1): 41-47
- Zhang CZ & Jin LF (1992) Preliminary studies on the bionomics of *Haplogonatopus japonicus*, *Natural Enemies of Insects*, **14** (2): 57-61
- Zhang ZQ (1991) Effects of draining paddy fields for control of white backed planthopper, *Insect Knowledge*, **28** (6): 321-325
- Zhu ZR, Cheng JA & Chen X (1993) Host preference and suitability of *Anagrus nilaparvatae*, *Acta Entomologica Sinica*, **14** (4): 430-437

Not accessed because of language - but look useful

- Abe Y, Koyama K, 1991. Embryonic development and selective oviposition of a dryinid wasp, *Haplogonatopus atratus* Esaki et Hashimoto (Hymenoptera: Dryinidae). *Japanese Journal of Applied Entomology and Zoology*, **35**(1):57-63. (Japanese)
- Chen YX (1987) Preliminary observations on habits of dryinid wasp of planthoppers in the east of Guizhou Province, *Insect Knowledge*, **24** (4): 237-239 (in Chinese – English abstract)
- Cheng JA & He JH, (1996) *Rice Insect Pest*. China Agricultural Press, Beijing, China (in Chinese).
- Cheng X & Xu G (1991) Comparison of population ecology between 2 mymarid egg parasitoids of rice planthoppers. *Acta Entomol. Sin*, **34**: 405-412 (in Chinese)
- Flint P, Magor JI, 1982. Bibliography and geographical index of literature on the brown planthopper, *Nilaparvata lugens* (Stal). Miscellaneous Report, Centre for Overseas Pest Research, No. 58:163 pp. (German)
- Jin X & Zhang C (1980) A preliminary study on the egg parasitoids of rice planthoppers, *Anagrus* spp., *Natural Enemies of Insects*, **2**: 51-57 (in Chinese)
- Luo XN, Zhuo WX, (1980) Studies on egg parasites of rice planthoppers *Anagrus* (I) *Entomol. Knowledge* **17**, 105-110 (in Chinese).
- Luo XN, Zhuo WX, (1980) Studies on egg parasites of rice planthoppers *Anagrus* (II) *Entomol. Knowledge* **17**, 147-151 (in Chinese).
- Luo XN, Zhuo WX, (1981) Studies on parasitoids of rice planthoppers *Anagrus* spp. Preference for hosts of three species A. spp. *Entomol. Knowledge* **18**, 3–6 (in Chinese).
- Luo XN, Zhuo WX, Chen, DR, (1981) Several kind of pesticides to rice plant hopper egg parasitic wasp—*Anagrus* spp.'s toxic observation. *Plant Prot. (China)* **5**, 24–25 (in Chinese).
- Luo XN, Zhuo WX, Chen DR, (1983) Bringing the function of egg-parasites on *Nilaparvata lugens* into full play, *Natural Enemies of Insects*, **5**: 61-63 (in Chinese)
- Chui SC (1978) The natural enemies of green leafhopper and brown planthopper, In: Ren-jong and Chiu (eds) *Diseases and insect pests of rice: ecology and epidemiology*, (in Chinese with English summary).
- Lin KS (1974) Notes on some natural enemies of *Nephotettix cinicticeps* (Uhler) and *Nilaparvata lugens* (Stål) in Taiwan. *Jour. Taiwan Agric. Res*, **23**: 91-115 (in Chinese with English summary).
- Lin G & Hu J (1985)
- Wang Y & Pang X (1986) Investigation of the host range *Anagrus nilaparvatae* (Hym. Myrmaridae), *Natural Enemies of Insects*, **8**: 225-229 (in Chinese)
- Xu GM & Cheng X (1990) Functional response to host density and mutual interference of *Anagrus nilaparvatae*, an egg parasitoid of the rice planthoppers, *Natural Enemies of Insects*, **12**: 51-55 (in Chinese)
- Xu HX & Ju RT (2003) Parasitization adaptability of *Anagrus optabilis* on *Nilaparvata lugens*, *Chinese Journal of Biological Control*, **19** (3): 136-138

Yuno K, Ehara S, 1982. Erythracid mites (Acarina, Erythracidae) parasitic on planthoppers. *Kontyu*, 50:344-345. (Japanese?)

Zhu ZR, Cheng JA, Chen X, 1993. Host preference and suitability of *Anagrus nilaparvatae*. *Acta Entomologica Sinica*, 36(4):430-437 (Chinese)